

# The Synapse

*The Medical Professionals' Network*

Lydgate, brain drain and the  
Maltese medical profession p10

Management of consumers' pharmaceutical  
waste in a pharmacy setting p16

Exclusive

**The Malta  
Foundation  
Programme -  
One year on**

p 6

**What is Doppler ultrasound?**

**Healing & Disease Reversal  
– The Series**

**What is on for Health Promotion  
and Disease Prevention  
Directorate for summer?**

**Meeting Carmen Joslin**



**OTHER INDICATIONS:**

- Treatment of GIO
- Male osteoporosis

**ACLASTA®**

# THE ONCE-YEARLY INFUSION OF POWERFUL OSTEOPROTECTION

## FOR POSTMENOPAUSAL OSTEOPOROSIS

- Significantly reduced 3-year risk of fractures at all key osteoporotic sites\*

**70%**

risk reduction in vertebral fracture†

**41%**

risk reduction in hip fracture†

**25%**

risk reduction in nonvertebral fracture\*\*\*

- A 15 minute, once-yearly infusion ensures yearlong compliance†
- Most adverse events were transient and mild to moderate‡
- Patient-preferred over weekly oral alendronate‡

\*Relative to placebo.

\*\*\*Nonvertebral fracture includes wrist, rib, arm, shoulder, or hip fracture; excludes finger, toe, or craniofacial fracture†

### ACLASTA® 5 MG (zoledronic acid) solution for infusion

**PRESENTATION:** Zoledronic acid. 100 mL solution bottle contains 5 mg zoledronic acid (anhydrous), corresponding to 5.330 mg zoledronic acid monohydrate.

**INDICATIONS:** Treatment of osteoporosis in post-menopausal women and men at increased risk of fracture, including those with a recent low-trauma hip fracture. Treatment of osteoporosis associated with long-term systemic glucocorticoid therapy in postmenopausal women and in men at increased risk of fracture. Treatment of Paget's disease of the bone. **DOSAGE AND ADMINISTRATION:** Osteoporosis: A single intravenous infusion of 5 mg Aclasta administered once a year. In patients with a recent low-trauma hip fracture, it is recommended to give the Aclasta infusion two or more weeks after hip fracture repair. Paget's Disease: A single intravenous infusion of 5 mg Aclasta. Specific re-treatment data are not available for Paget's disease. Aclasta is administered via a vented infusion line and given at a constant infusion rate. The infusion time must not be less than 15 minutes. Adequate calcium and vitamin D are recommended in association with Aclasta administration. In patients with recent low-trauma hip fracture a loading dose of 50,000 to 125,000 IU of Vitamin D is recommended prior to the first Aclasta infusion. No dose adjustment in patients with creatinine clearance  $\geq 35$  mL/min, or in patients with hepatic impairment, or in elderly patients. The safety and efficacy of Aclasta in children and adolescents below 18 years of age has not been established. **CONTRAINDICATIONS:** Hypersensitivity to zoledronic acid or to any of the excipients or to any bisphosphonate; hypocalcaemia; pregnancy; lactation. **PRECAUTIONS AND WARNINGS:** Serum creatinine should be measured before each Aclasta dose. Aclasta should not be used in patients with creatinine clearance  $< 35$  mL/min. Transient increase in serum creatinine may be greater in patients with underlying impaired renal function. Monitoring of serum creatinine should be considered in at-risk patients. Patients must be appropriately hydrated prior to administration of Aclasta, especially important for the elderly and for patients receiving diuretic therapy. Use with caution in conjunction with medicinal products that can impact renal function. A single dose of Aclasta should not exceed 5mg and the duration of infusion should be at least 15 minutes. Pre-existing hypocalcaemia and other disturbances of mineral metabolism must be treated by adequate intake of calcium and vitamin D before initiating therapy with Aclasta. It is strongly advised that patients with Paget's disease receive supplemental calcium and vitamin D. Measurement of serum calcium before infusion is recommended for patients with Paget's disease. Severe and occasionally incapacitating bone, joint and/or muscle pain have been infrequently reported with bisphosphonate therapy. A patient being treated with Zometa should not be treated with Aclasta. As a precaution against osteonecrosis of the jaw (ONJ) a dental examination with appropriate preventive dentistry should be considered prior to treatment in patients with concomitant risk factors (e.g. cancer, chemotherapy, corticosteroids, poor oral hygiene). While on treatment, these patients should avoid invasive dental procedures if possible. For patients requiring dental procedures, there are no data available to suggest whether discontinuation of bisphosphonate treatment reduces the risk of osteonecrosis of the jaw. Aclasta is not recommended in women of childbearing potential. **INTERACTIONS:** Specific drug-drug interaction studies have not been conducted with zoledronic acid. Caution is recommended when Aclasta is used concomitantly with drugs that can significantly impact renal function, such as aminoglycosides and diuretics that can cause dehydration. In patients with renal impairment, the systemic exposure to concomitant medicinal products that are primarily excreted via the kidney may increase. **ADVERSE REACTIONS:** The incidence of adverse reactions (e.g. fever, myalgia, flu-like symptoms, arthralgia and headache) are greatest with the first infusion and decrease markedly with subsequent infusions. The majority of these reactions occur within the first three days and were mild to moderate and resolved within three days of the event onset. The incidence of these adverse reactions can be reduced with the administration of paracetamol or ibuprofen shortly following Aclasta administration. Very common: Fever. Common: Flu-like symptoms, chills, fatigue, pain, asthenia, malaise, arthralgia, myalgia, bone pain, back pain, pain in extremity, vomiting, nausea, headache, dizziness, atrial fibrillation, hypocalcaemia†, ocular hyperaemia, diarrhoea, increased C-reactive protein, infusion site reactions. Uncommon: Hypertension, flushing, palpitations and others. Not known: Scleritis, orbital inflammation, hypotension, renal impairment, osteonecrosis of the jaw, dehydration secondary to post dose symptoms, hypersensitivity reactions † *Common in Paget's disease only*. Please refer to SmPC for a full list of adverse events. **PACK SIZE:** Aclasta is supplied in packs containing one 100ml bottle. **LEGAL CATEGORY:** POM. **MARKETING AUTHORISATION NUMBER:** EU/1/05/308/001. **MARKETING AUTHORISATION HOLDER:** Novartis Europharm Limited, Wimblehurst Road, Horsham, West Sussex, RH12 5AB, United Kingdom. Consult full Summary of Product Characteristics (SmPC) before prescribing. Full prescribing information is available upon request from: Novartis Pharma, P.O. Box 124, Valletta VLT 1000, Malta. Tel +356 22983217. (vs n 2010-M-001 ACL 18-05-2010)

**References:** 1. Aclasta SmPC. Novartis Europharm Ltd. 2. Black DM, Delmas PD, Eastell R, et al; for the HORIZON Pivotal Fracture Trial. Once-yearly zoledronic acid for treatment of postmenopausal osteoporosis. *N Engl J Med.* 2007;356:1809-1822. 3. Saag K, Lindsay R, Kriegman A, Beamer E, Zhou W. A single zoledronic acid infusion reduces bone resorption markers more rapidly than weekly oral alendronate in postmenopausal women with low bone mineral density. *Bone.* 2007;40:1238-1243. 4. McClung M, Recker R, Miller P, et al. Intravenous zoledronic acid 5mg in the treatment of postmenopausal women with low bone density previously treated with alendronate. *Bone.* 2007;41:122-128.




**Aclasta®**  
zoledronic acid 5 mg  
solution for infusion  
*One Infusion. Yearlong Osteoprotection.*

# Of Social Corporate Responsibilities

These last 2 months I have been suffering from bilateral arm nocturnal paresthesia. Probably it is carpal tunnel syndrome, they have told me, but I am awaiting for an EMG examination to check out matters. Not exactly what I have wished for in my New Year Resolution, so I took refuge yet again in one of Paolo Coelho books... The devil and Mrs Pym. For those of you who never heard of this Brazilian songwriter-turned-author (he also wrote The Alchemist, recently also translated in Maltese), they are missing much. His writings are a man's spiritual journey... the particular novel which I am currently reading explores the timeless struggle between good and evil, and brings to our everyday dilemmas a fresh perspective: how we can be incentivised to master the fear that prevents us from following our dreams, from being different, from truly living.

Indeed as I read this book, I reflect, as I always do when I read Coelho's books. Indeed, as years are beginning to etch their markings on me, I am becoming more aware of how stupid we can be when our pursuit of money turns from simply a need to a want. We start to forget what we are and what being a human being really means.

We may start to neglect our spouses, our children, our ethical standards... possibly because of money and lust. Indeed I strongly believe that we, as healthcare professionals, have an obligation towards others, not simply by offering a service for which we are paid for but also by offering our time and resources free of charge to help others and to pursue altruism. I admire mostly those healthcare professionals who involve themselves on a voluntary basis in patient support groups (obviously excluding those who do so to increase their clientele).

A recent group worth mentioning is the Professionals against Embryo Freezing. Obviously not all of you may agree with their agenda but everyone will agree that this group is indeed extraordinary. To put it simply, various professionals including doctors, pharmacists and dentists have joined their grey matter together, forming a group over a very short time (2 weeks) to pursue an ideal. And this is what I admire most of them. Not only because of their impeccable division of labour coordinated by Dr Miriam Sciberras. But because seeing this inter and intra-professional cooperation rekindles my hope that there are still healthcare

professionals who indeed are willing to contribute part of their time free of charge to pursue altruism.

Hopefully we will see more of these professional groups pursuing more ideals... those very ideals which are the very foundations of our noble and respected medical professional including beneficence, non-maleficence and justice...

Over the past years The Synapse has offered free advertising space to various patient support groups. The latter have submitted articles on their terms of reference with some also advertising conferences and other fund raising events. However a recent tool introduced by The Synapse Website portal should be of particular benefit to them (and to other healthcare professionals practicing in other fields), namely these patient groups may directly reach their peers by uploading their events online - [www.thesynapse.net/events/list.asp](http://www.thesynapse.net/events/list.asp). And what is even more convenient is that this is also free of charge. §

*Ian C Ellul*

Ian C Ellul





**DaVinci**  
HOSPITAL

**Call 21 491 200**



# a turning point for weight loss

**real evidence** – **alli** is orlistat 60 mg, the first and only EU-licensed non-prescription weight loss treatment.

**real help** – **alli** combines a capsule and a support programme to help users lose 50% more weight than by diet alone!<sup>1</sup>

**real benefits** – **alli** brings positive change to customers and the opportunity for you to recommend with confidence.



orlistat

**start the conversation**  
find out more at [www.alli.com.mt](http://www.alli.com.mt)



**Product Information.** **alli** 60 mg hard capsules (orlistat). **Indication:** Weight loss in adults BMI ≥ 28. **Dosage:** Adults (18 or over): One capsule with each of three main meals. Max. 3 caps/day for up to 6 months. Use with mildly hypocaloric, lower-fat diet. If no weight loss within 12 weeks, refer to doctor or pharmacist. Diet and exercise should start prior to treatment. **Contraindications:** Hypersensitivity to ingredients; concurrent treatment with oral anticoagulants or ciclosporin; chronic malabsorption syndrome; cholestasis; pregnancy; breast-feeding. **Special warnings and precautions:** Talk to doctor before starting to take **alli** if taking amiodarone, a medicine for diabetes, epilepsy or hypothyroidism, or if patient has kidney disease. If taking a medicine for hypertension or hypercholesterolaemia, talk to doctor or pharmacist when taking **alli**. Risk of gastrointestinal (GI) symptoms increases with fat consumption.

Take multivitamin at bedtime. See doctor if rectal bleeding occurs. Oral contraceptive efficacy may be reduced if severe diarrhoea occurs; use additional contraception. **Drug interactions:** Ciclosporin; oral anticoagulants; levothyroxine; antiepileptic medication; oral contraception; fat soluble vitamins; acarbose; amiodarone. **Pregnancy and lactation:** Do not use during pregnancy or lactation. **Side effects:** See SPC for full details. Predominantly GI e.g. oily stools, urgency; usually mild and transient, risk reduced by low fat consumption. Diverticulitis; pancreatitis; mild rectal bleeding; hepatitis; cholelithiasis; abnormal liver enzymes; anxiety; hypersensitivity reactions including anaphylaxis, bronchospasm, angioedema, pruritus, rash, and urticaria; bullous eruption. **Legal category:** Non-prescription. Marketing Authorisation Holder: Glaxo Group Limited, Greenford, Middlesex, UB6 0NN. **MA Number:** EU/1/07/401/007 - 010. Pack size 84s. **Last revised:** March 2010.

**alli** is a registered trademark of the GlaxoSmithKline group of companies



**Dr Charmaine Gauci** MD MSc Dip(Fit&Nut) PhD FRSPH FFPH is the Director of the Health Promotion and Disease Prevention Directorate. She is a senior lecturer with the University of Malta and delivers lectures in the field of public health with special interest in Epidemiology and Communicable Diseases. She is active in the field of public health and is currently also the President of the Malta Association of Public Health Medicine.



**Karen Attard** is a final year Pharmacy student. She has attended the Erasmus program in Germany where she worked on a research project on Kollidon® microparticles as sustained release preparations. The research was carried out under the supervision of Professor L. Azzopardi, Professor A. Serracino-Inglott and Dr. M. Zarb-Adami from the Department of Pharmacy.



**Professor Kevin Cassar** MD(Malta) MMed(Dundee) FRCS(Edin) MD(Aberdeen) FRCS(Intercoll) is Consultant Vascular Surgeon at Mater Dei Hospital and Associate Professor at the University of Malta. He is the author of several peer reviewed articles and book chapters including the Peripheral Arterial Disease section of the BMJ Clinical Evidence. The co-author of the article is Dr Tonio Piscopo.



**Professor Albert Cilia-Vincenti** MD FRCPath is chairman of the Academy of Nutritional Medicine of UK, and a private surgical pathologist in Malta. He is a former pathology teacher at London and Malta universities, and pathology services director to the British and Maltese health services. He trained at London's Royal Marsden, Royal Free, St George's, Charing Cross and The Middlesex hospitals.



**Professor Victor Grech** MD PhD is a consultant paediatrician with a special interest in paediatric cardiology. He is also the creator and editor-in-chief of the journal Images in Paediatric Cardiology ([www.impaedcard.com](http://www.impaedcard.com)).



**Dr Pierre Vassallo** MD PhD FACA Artz für Radiologie specialised in radiology at the Institute of Clinical Radiology at the University of Muenster, Germany and the Memorial Sloan-Kettering Cancer Center, New York, US. He is currently Consultant Radiologist and Managing Director at DaVinci Hospital, Malta.

# contents

- 06 The Malta Foundation Programme: one year on – Part II
- 09 What is on for the Health Promotion and Disease Prevention Directorate for summer?
- 10 Lydgate, brain drain and the Maltese medical profession
- 13 Healing & Disease Reversal THE SERIES
- 15 Network announcements
- 18 Management of consumers' pharmaceutical waste in a pharmacy setting
- 20 Meeting Dr Carmen Joslin
- 25 What is Doppler ultrasound?


Published by Medical Portals Ltd.  
The Professional Services Centre  
Guzi Cutajar Street  
Dingli, Malta  
Email: [editor@thesynapse.net](mailto:editor@thesynapse.net)  
Web: [www.thesynapse.net](http://www.thesynapse.net)

Editor: Wilfred Galea  
Scientific Editor: Ian C Ellul  
Administration Manager: Carmen Cachia

Production: Outlook Coop  
Printing: Europrint Ltd

The opinions expressed in this publication are those of the respective authors and do not necessarily reflect the opinions of the editors or the institutions with which the author is affiliated unless this is clearly specified.

**Advertising policy:** Advertisers are liable for contents of any of the advertisements. The advertisers shall indemnify and hold harmless Medical Portals Ltd against and from any and all claims, damages, liabilities, cost and expenses whatsoever, including counsel fees, arising from the content of any of their advertisements. Medical Portals Ltd disclaims any responsibility or liability for non-compliance of advertising artwork to regulatory units.



**COVER:** Capparispinus (Kapparis, Caper)

#### Medicinal uses:

A perennial shrub growing on rocky soil, the buds of which are a common ingredient in Mediterranean cuisine. As a herbal tea or a compress made from its roots or young shoots, caper has been used medicinally to counter the pain of sciatica or gout.

Reference: Lanfranco G. Hxejex Mediċinali u oħrajn fil-gżejjer Maltin. Media Centre Print; Malta. 1993

Photography: Guido Bonett ARPS AMPS

# The Malta Foundation Programme: One year on - Part II

CONTINUED FROM ISSUE 2/11

### 3. Assessment

Assessment of trainees is another major change brought about by the foundation programme. Whereas prior to the introduction of the programme, trainees were given full registration with the Medical Council of Malta without any formal assessment, trainees are now assessed repeatedly during the two years using different assessment tools and by several different assessors. In order to implement these changes over 250 assessors received training in the assessment tools by the Malta Foundation School. The tools include mini-Clinical Examination (mini-CEX) which assesses the trainee's interaction with a patient, Direct Observation of Procedural Skills (DOPS) which assesses the trainee's competence in performing a procedure, Case-Based discussion (CBD) which assesses the trainee's management, documentation and professionalism, and multisource feedback (MSF) which is a 360 degree assessment by at least 10 individuals with whom the trainee works.

Trainees are obliged to complete at least 20 of these assessments during the year. Introducing this intensity of assessment was of course a major challenge and the considerable number of assessments raised some concerns about whether this was feasible, particularly as no formal assessments of trainees had been carried out in the past. The first year has shown us that our misgivings were unfounded. For 86 trainees, between July 2009 and July 2010, no less than 2317 MSFs, 516 CBDs, 1117 DOPS, and 501 mini-CEXs were carried out. These are huge numbers of assessments and achieving these numbers was only possible through the cooperation and efforts of a significant number of people, not least by the Consultant body. Indeed our records indicate that 72% of all CBDs and 23% of all mini-CEXs were carried out by consultants. The UK's

Postgraduate Medical Education Training Board (PMETB) survey of 2009 reported that in the UK only 46% of CBDs are carried out by consultants. This indicates that consultant involvement in the assessment of our trainees is far higher than it is in the UK. It also indicates that consultants are making a very major contribution not only to the training of foundation doctors but also to their assessments.



THE ROLE OF THE TEAM  
SET UP LOCALLY  
IS TO ASSESS TRAINEES  
WHO ARE EXPERIENCING  
DIFFICULTIES TO TRY AND  
ESTABLISH THE CAUSE  
OF THE PROBLEMS  
AND TO PROVIDE  
THE SUPPORT REQUIRED

### 4. Trainee Support

A trainee support team (TST) has been set up within the Malta Foundation School under the direction of Dr Etienne Muscat. The aim of the TST is to provide junior doctors enrolled in the Foundation Programme with access to a system that can assist them with issues that are hindering them in their training and, by extrapolation, in their expected maturation into

independent, reliable and safe doctors. The concerns raised with the TST may include deficiencies of knowledge, clinical skills, professionalism, behaviour, substance abuse, and mental health and may be raised by the doctor or supervisor. Prior to the setting up of this team, Dr Etienne Muscat visited the Trainee Support Unit at East Midlands Foundation Schools in the UK to witness first hand the functioning of the unit. The role of the team set up locally is to assess trainees who are experiencing difficulties to try and establish the cause of the problems and to provide the support required. During the first year of the foundation programme 11 trainees were assessed, 9 at FY1 and 2 at FY2 level. Interventions in these trainees included referral for clinical psychology evaluation, counselling therapy service, referral for independent psychiatric board evaluation, and referral for careers counselling.

### 5. Careers Advice

Dr Pierre Ellul is responsible for provision of careers advice to foundation doctors. The role of Careers advice is to organise careers workshops, to support the development of taster weeks, to provide doctors with careers information, and to hold careers planning meetings with trainees as required. Last year all FY2 trainees were offered Windmills Careers Day. Dr Pierre Ellul had received training in delivery of this careers workshop and delivered the workshops together with UK trainers. The feedback from our trainees on this workshop was excellent.

### 6. Feedback

Throughout the year the trainees' feedback on various aspects of their training has been sought. This includes feedback about the lecture programme as well as their rotations. End-of-post questionnaires are conducted

**Table 1:** Number of doctors recruited into national health service

	2007	2008	2009	2010	2011
UOM final year student numbers	55	53	59	57	76
Final year students joining health service	35	38	47	48	68
% Joining health service	63.6	71.4	79.7	84.2	90
Total number recruited	35	38	50	54	88
<b>% of final year numbers recruited</b>	<b>63.6</b>	<b>71.4</b>	<b>84.7</b>	<b>94.7</b>	<b>115.7</b>

anonymously using an on-line survey. 94% of trainees stated that they were satisfied or highly satisfied with the clinical skills acquired during their assignment (Figure 1). 95% reported that they frequently, regularly or sometimes received feedback (Figure 2) which the majority (93%) found useful (Figure 3). Feedback from trainees also indicates that only a small minority of trainees were dissatisfied or very dissatisfied with the training provided

by their clinical supervisor (15.9%) with the majority being very satisfied or satisfied.

### 7. Recruitment

The second objective of the foundation programme was to improve the retention rate of medical graduates within the service. In 2007, only 63.6% of graduates from the University of Malta were recruited to the local health service. The figure in 2008 was slightly better at 71.4%. The introduction of the foundation programme resulted not only in an improvement in retention of local graduates but also attracted foreign graduates to join the programme. In 2010, 84.2% of University of Malta graduates joined the Malta foundation programme together with 8 foreign graduates. As a result the total number of doctors recruited in 2010 reached almost 95% of the number of students graduating from the University of Malta (Table 1). The application process for entry into the Foundation programme in July 2011 has been completed. For the first time the number of doctors recruited

(88) exceeds the total number of final year medical students at the University of Malta (76). This indicates that this second objective has definitely been reached.


### Conclusions

The setting up of the foundation programme has resulted in major changes in the training of 'houseofficers'. It has provided a structured programme with good quality training, supervision and assessment. The number of doctors recruited to the service has increased steadily and is expected to continue to increase. This will not only improve the service but also the quality of training through increased educational opportunities and better distribution of workload.


The results obtained however do not allow for complacency. There are major challenges ahead, not least the impending full implementation of the European Working Time Directive, increasing numbers of medical students at the University of Malta, increasing competition for posts from other EU nationals, increased demand on educational resources, and of course need for more basic specialist training posts for the increasing number of doctors.

The successful introduction of the Foundation programme in Malta is an example of how co-operation between all stakeholders together with the hard work of clinical and administrative staff can lead to major improvements in training standards and the rapid acquisition of a common goal. §


**Figure 1:** Responses to question "Are you satisfied with the clinical skills acquired during this assignment?"



**Figure 2:** Responses to "Did you receive feedback about your performance during this assignment?"



**Figure 3:** Responses to "Was the feedback you received useful?"



A REVOLUTIONARY  
TREATMENT FOR  
WOMEN WITH  
POSTMENOPAUSAL  
OSTEOPOROSIS

## A NEW FORCE AGAINST FRACTURE

STOPPING OSTEOCLASTS BEFORE THEY REACH THE BONE

Introducing the first and only  
RANK Ligand inhibitor  
that works throughout the  
skeleton to protect women with  
postmenopausal osteoporosis.<sup>1</sup>

 **prolia**<sup>®</sup>  
denosumab

A FORCE AGAINST FRACTURE

**Reference:** 1. Prolia<sup>®</sup> Summary of Product Characteristics, 2010.

### Prolia<sup>®</sup> (denosumab) Brief Prescribing Information

Please refer to the SmPC (Summary of Product Characteristics) before prescribing Prolia<sup>®</sup>. **Pharmaceutical Form:** 1 ml solution for injection presented in pre-filled syringe containing 60 mg of denosumab. Contains sorbitol [E420]. **Indications:** Treatment of osteoporosis in postmenopausal women at increased risk of fractures. Prolia<sup>®</sup> significantly reduces the risk of vertebral, non-vertebral and hip fractures. **Dosage and Administration:** Single subcutaneous injection of Prolia<sup>®</sup> 60 mg is given once every 6 months. No dose adjustment for renal impaired patients. Patients must be supplemented with calcium and vitamin D. Prolia<sup>®</sup> is not recommended in paediatric patients (age < 18). **Contraindications:** Hypocalcaemia. Hypersensitivity to the active substance or any of the excipients. **Special warnings and precautions for use:** Hypocalcaemia must be corrected by adequate intake of calcium and vitamin D before initiating therapy. Patients with severe renal impairment or receiving dialysis are at greater risk of hypocalcaemia. Clinical monitoring of calcium levels is recommended for patients predisposed to hypocalcaemia. Patients receiving Prolia<sup>®</sup> may develop skin infections (predominantly cellulitis) leading to hospitalisation and should contact

a healthcare professional immediately if they develop signs or symptoms of cellulitis. Osteonecrosis of the jaw (ONJ) has been reported with denosumab and with bisphosphonates. ONJ has been reported rarely with Prolia<sup>®</sup> 60 mg every 6 months. A dental examination should be considered prior to treatment with Prolia<sup>®</sup> in patients with concomitant risk factors (refer to SmPC). While on treatment, these patients should avoid invasive dental procedures if possible. Good oral hygiene practices should be maintained during treatment with Prolia<sup>®</sup>. The needle cover of the syringe contains dry natural rubber (latex derivative), which may cause allergic reactions. Patients with rare hereditary problems of fructose intolerance should not use Prolia<sup>®</sup>. **Interactions:** No interaction studies have been performed. The potential for pharmacodynamic interactions with hormone replacement therapy (HRT) is considered to be low. **Pregnancy and lactation:** Prolia<sup>®</sup> is not recommended for use in pregnant women. A risk/benefit decision should be made in patients who are breast feeding. It is unknown whether Prolia<sup>®</sup> is excreted in human milk. No data are available on the effect of Prolia<sup>®</sup> on human fertility. **Undesirable effects:** Adverse reactions reported in placebo-controlled clinical studies

in women with postmenopausal osteoporosis and breast or prostate cancer patients receiving hormone ablation: **Common** (> 1/100, < 1/10) Urinary tract infection, Upper respiratory tract infection, Sciatica, Cataracts, Constipation, Rash, Pain in extremity; **Uncommon** (> 1/1,000, < 1/100) Diverticulitis, Cellulitis, Ear infection, Eczema; **Very Rare** (< 1/10,000) Hypocalcaemia. In the osteoporosis clinical program ONJ has been reported rarely with Prolia<sup>®</sup>. Please consult the SmPC for a full description of side effects. **Pharmaceutical Precautions:** Do not mix with other medicinal products. Store in a refrigerator (2°C–8°C). Do not freeze. Keep the pre-filled syringe in the outer carton in order to protect from light. Do not shake excessively. Prolia<sup>®</sup> may be stored at room temperature (up to 25°C) for up to 30 days in the original container. Once removed from the refrigerator use within these 30 days. **Marketing authorisation holder:** Amgen Europe B.V., Minervum 7061, NL-4817 ZK Breda, The Netherlands. Further information is available from the SmPC. Date of PI preparation: May 2010. Adverse events should be reported. **Legal Category:** Medicinal product subject to medical prescription. **Marketing authorisation number:** EU/1/10/618/003.

To learn more, visit: [www.prolia-international.com](http://www.prolia-international.com)

**AMGEN**<sup>®</sup>

 **gsk**  
GlaxoSmithKline

© 2010 Amgen, Zug, Switzerland.  
All rights reserved.

NP-AMG-547-2009  
04.2010





# What is on for the Health Promotion and Disease Prevention Directorate for summer?

CHARMAINE GAUCI

**E**xcessive sun exposure can cause many different complications. Over-exposure to the sun can cause a sunburn and permanent damage to the skin. This can lead to the development of skin cancer. Adolescents and young children are at a greater risk of skin damage due to over-exposure of the sun. Other effects of sun exposure include actinic keratosis and spots. It is also evident that it can result in wrinkles and premature ageing of the skin.

The Health Promotion and Disease Prevention Directorate focuses on protection from sun exposure. Emphasis is made during summer however protection is important all year round in every setting.


Some tips for prevention recommended by the World Health Organisation include:

- **Limit time in the mid-day sun**  
The sun's UV rays are the strongest between 10 a.m. and 4 p.m. One should limit exposure to the sun during these hours.
- **Watch for the UV index**  
This important resource helps you plan your outdoor activities in ways that prevent over-exposure to the sun's rays. While you should always take precautions against over-exposure, take special care to adopt sun safety practices when the UV Index predicts exposure levels of moderate or above levels.

- **Use shade wisely**  
Seek shade when UV rays are the most intense, but keep in mind that shade structures such as trees, umbrellas or canopies do not offer complete sun protection. Remember the shadow rule: "Watch your shadow – Short shadow, seek shade!"
- **Wear protective clothing**  
A hat with a wide brim offers good sun protection for your eyes, ears, face, and the back or your neck. Sunglasses that provide 99 to 100 percent UV-A and UV-B protection will greatly reduce eye damage from sun exposure. Tightly woven, loose-fitting clothes will provide additional protection from the sun.
- **Use sunscreen**  
Apply a broad-spectrum sunscreen of SPF 15+ liberally and re-apply every two hours, or after working, swimming, playing or exercising outdoors.
- **Avoid sunlamps and tanning parlours**  
Sunbeds damage the skin and unprotected eyes and are best avoided.

Secondary prevention in raising awareness about identification of early stages of skin cancer is important. Melanoma is the most serious form of skin cancer in terms of mortality. It

may develop anywhere on the skin, and within a pre-existing mole in 30 to 50% of cases. The commonest sites are the trunk in men, and the legs and trunk in women however it can develop spontaneously. Hence clinicians should:

- **Be aware that fair-skinned men and women aged 65 and older, and people with atypical moles, and people with more than 50 moles, are at greater risk for developing melanoma.**
- **Remain alert for skin abnormalities when conducting physical examinations for other purposes.** The A-B-C-D-E rule can be useful in making an early diagnosis ie if a mole is asymmetrical (A); has no distinct border (B); if its colour changes (C); if it is larger than 5mm in diameter (D); evolution (E), such as changes in color, shape, size, elevation, skin surface, and symptoms such as itching or bleeding of a lesion are a hallmark sign of malignancy. 



VICTOR GRECH

# Lydgate, brain drain and the Maltese medical profession

Globalisation causes the migration of all types of workers, including healthcare professionals, who may opt to seek richer pastures in the form of better paid and pensionable positions in countries other than their own. Dr Tertius Lydgate in Eliot's 19<sup>th</sup> century novel *Middlemarch*, while starting out with lofty ideals, rapidly became mired in financial difficulties and was constrained to seek more lucrative pastures outside Middlemarch. Maltese doctors are also poorly paid, and unless the local authorities ameliorate working conditions, the ongoing medical brain drain will worsen with catastrophic consequences for the local NHS.

Globalisation is fashioning a workforce resettlement that also encompasses the medical profession, and Maltese doctors are no exception to these migrations. We shall briefly review Tertius Lydgate, a medical character in George Eliot's *Middlemarch*, a protagonist who even in the 19<sup>th</sup> century, clearly demonstrated the possibilities and implications of such migrations.<sup>1</sup>

However, we shall first briefly review the background to doctors in English literature up to Eliot's period. Medics in English literature were initially viewed with irony and treated with disdain, and one of the earliest such characters is Chaucer's 'Doctour of Phisik' in his 14<sup>th</sup> century *Canterbury Tales*, a knowledgeable man who is depicted as avaricious and in league with apothecaries who overcharge patients for cheap medicines. By the time we reach William Shakespeare's period (1564-1616), attitudes had somewhat improved, despite the chaotic mess that medical regulatory affairs found themselves in, example, quacks and empirics practiced widely and unsafely, the distinction between physicians and surgeons was blurred, the Physicians Act of 1540 defined medicine as also encompassing surgery, giving the physicians the right to operate, and a Guild of Surgeons had existed since 1369 while an older Guild of Barbers included members who practiced either exclusively as surgeons and tooth-drawers or only shaving and hairdressing. Moreover, apothecaries were members of the Grocers' Company. This tripartite system of medical practice in Tudor England was evidently highly confusing and a 1512 Act of Parliament had acknowledged that a vast number of untrained individuals were practicing medicine, stating that none could practice medicine in London or within a seven-mile radius unless graduates of Oxford or Cambridge or upon having passed an examination by the Bishop of London or the Dean of St Paul's Church.<sup>2</sup>



Overall, Shakespeare appears to have thought highly of physicians who are mentioned as learned doctors, while viewing surgeons with a jaundiced eye, with, for example Dick Surgeon in *Twelfth Night* discovered in a state of drunken stupor when most needed. By the beginning of the 19<sup>th</sup> century, the traditional medical responsibilities of physicians, surgeons, and apothecaries were being replaced by new and well-schooled practitioners who could perform all of these roles. George Eliot's fictional Dr. Lydgate is one of these novel medical practitioners, and is one of *Middlemarch*'s protagonists. His dilemmas, at work and at home, are used to illuminate social, psychological, professional and moral problems. Lydgate is first mentioned well into the novel, in chapter 10, and although an aristocrat, is poor, ambitious and somewhat arrogant.

The noble doctor had deliberately kept away from London's big city intrigues, jealousies, social jockeying and the jostle for celebrity status. He starts off with lofty ideals that include the running of a fever hospital in rural Middlemarch for free and the performance of medical research into the causes of fever and other illnesses. This causes the town's medical establishment to treat him with jealousy and suspicion and Lydgate's practice consequently develops very slowly. While Lydgate is a moral man, like us all, he suffers from a number of small prejudices and moral failings that are related to the necessity of balancing self-interest with that of other individuals and patients.<sup>3</sup>

Lydgate's downfall occurs when he falls in love with, and marries, a pretty and highly decorative but thoroughly impractical woman who is used to a rich lifestyle. She quickly depletes his savings and goes on to mire him in dire financial straits, dragging him progressively deeper into debt. Lydgate finds himself eventually implicated in malpractice during the death throes of one of his patients due to a potential conflict of interest that would have yielded him substantial financial benefit, releasing him from debt.

The eventual victory of society, personified by his wife Rosamond, over Lydgate's integrity is inescapable. Lydgate is forced to seek more lucrative pastures than Middlemarch among London's high society, abandoning his high ideals; a perfect Aristotelian tragic hero.<sup>4</sup>

Modern doctors are highly trained professionals, and expect to live reasonably comfortably. Furthermore, the inevitable contacts with peers in private practice, and fêting by drug and medical companies, suggests certain expectations of medical life. Pressures on the home front are equally unavoidable, with soaring property prices, and costs of maintaining a household with dependants that may comprise a wife and several children, including the latter's schooling costs.

All of these are exacerbated by the poor Maltese NHS pay scale. This situation has come about partly because the Maltese socialist administrations in the 1970s and 1980s regarded China, North Korea and Libya as ideal role-models, hence civil service pay scales were reduced and percentage wage increases were replaced by cost of living wage increases across the board, significantly lowering salaries and pensions of some of the best trained professionals in the country. Unfortunately, despite many years of non-socialist administration, pay scales have often failed to retain brain power that is highly appreciated and even actively head-hunted by medical agencies.<sup>5</sup> This also results in a significant financial loss to the country since the local authorities themselves have estimated that a medical student's costs at least Euro 95,000 to train in tertiary education expenditures alone.<sup>5</sup>

Also in UK NHS style, Maltese NHS salaries have traditionally been supplemented by private practice, and therefore specialities which attract little or no private practice tend to have difficulty recruiting trainees and specialists who prefer to move overseas in far better salaried and pensioned posts.<sup>6</sup>

Several solutions are available, and some are inherently unacceptable to the medical profession, including a lowering of future prospects that may, perhaps, be inculcated at the training level, a disingenuous expectation. First rate professionals cannot be expected for work for second rate salaries and the choices are stark. The danger of such deals was aptly highlighted by an eminent local retired pathology and ethics professor who pointed out the possibility that Maltese doctors may decide to take it upon themselves to decide what fraction of the 40-hour NHS week amounts to their salary, since an employee who believes he or she is being defrauded by one's employer is not morally wrong to diminish the work commitment to that employer, and that this would not be an unethical stance.<sup>7</sup>

This is probably more acceptable to individual doctors than the drastic lowering of one's standard of living or running up ruinous debts. The final alternative is emigration as clearly shown by Balzan et al. with 50% of medical graduates leaving the island to continue studying or work abroad but only 7.5% return.<sup>8</sup> The current Maltese administration has accepted the reality of this brain drain,<sup>5</sup> but the response to train much larger numbers of doctors is hardly likely to solve the brain drain problem as this will further exacerbate the local brain drain and contribute to the influx of well trained doctors to richer countries, particularly the United Kingdom (English is Malta's second language).

The Medical Association of Malta has proposed a three point plan to the government in order to encourage local doctors to remain and practice: an increase in basic salary in order to offset loss of income due to the introduction of the European working time directive,<sup>6</sup> the implementation of structured postgraduate training programmes in as many specialties as can be supported by the local infrastructure and a gradual expansion in the consultant grade in order to improve junior doctors' long term career prospects.

Middlemarch's Lydgate died when he was only fifty, leaving his wife and children provided for by a heavy life insurance. He had gained an excellent practice, alternating, between London and a Continental bathing-place by season and wrote a treatise on gout, a disease of the wealthy. And while he regarded himself as a failure as he failed to do what he had set out to do, Lydgate, ironically, was considered a successful man.

There is no easy solution to the problem of good doctors leaving state and academic service in poorer countries in favour of better remunerated employment with more favourable career prospects in both European and non-European countries.<sup>9</sup> The problem must be adequately, competently and urgently addressed by the local health authorities, by schemes such as the consolidation of the current Foundation program along with other encouragements and inducements that will persuade Maltese doctors who go abroad to train to return to their homeland. S

#### Acknowledgements

Drs. Ivan Callus and Claire Thake for their help and support in my ongoing work in English Literature in general.

#### References

1. George Eliot. *Middlemarch, a Study of Provincial Life*, London: William Blackwood and Sons, 1871.
2. K.T. Fu. The healing hand in literature: Shakespeare and surgery, *Hong Kong Med J*; 1998; 4:77-88.
3. A.T. Westwood. Lifestyle, Lydgate and the late twentieth century. *J R Soc Med*. 1997; 90:42-3.
4. D. Daiches. George Eliot's Dr Lydgate. *Proc R Soc Med*. 1971; 64:723-4.
5. F. Galea-Debono. Time for major investment in cancer equipment, *Times of Malta*; 2007.
6. D. P. Sandhu. Current dilemmas in overseas doctors' training. *Postgrad Med J*. 2005; 81:79-82.
7. A. Cilia-Vincenti. Medical brain drain (Talking Point). *Times of Malta*; 2006.
8. J. Ameen. Half of graduate doctors leave to work abroad – MAM study, *Malta Independent*; 2007.
9. I. Vlad. European views about accession. *BMJ Career Focus*. 2004; 328:99.

THROUGH DIRECT RENIN INHIBITION

THE FIRST SINGLE-PILL

COMBINATION WITH

**RASILEZ®**

**DRIVE POWER  
THAT LASTS**



**Rasilez HCT.**

aliskiren/hydrochlorothiazide

**DRIVE POWER THAT LASTS**

**RASILEZ HCT®: THE BENEFITS OF RASILEZ®  
WITH EVEN GREATER BP EFFICACY<sup>1,2</sup>**

**RASILEZ HCT® (aliskiren/hydrochlorothiazide) film coated tablets**

**PRESENTATION:** Each film coated tablet contains 300 mg aliskiren (a renin inhibitor) and 12.5 mg hydrochlorothiazide (a thiazide diuretic), or 300 mg aliskiren and 25 mg hydrochlorothiazide. **INDICATIONS:** Treatment of essential hypertension. Indicated in patients whose blood pressure is not adequately controlled on aliskiren or hydrochlorothiazide used alone. Indicated as substitution therapy in patients adequately controlled with aliskiren and hydrochlorothiazide, given concurrently, at the same dose levels as in the combination. **DOSEAGE:** One tablet of Rasilez HCT 300/12.5 mg or 300/25 mg daily with meals. No dose adjustment needed for renal impairment, mild to moderate hepatic impairment and elderly. **CONTRAINDICATIONS:** • known hypersensitivity to the components of this product or to sulfonamide-derived substances • history of angioedema with aliskiren • Second and third trimesters of pregnancy • breast-feeding • severe hepatic impairment • severe renal impairment (creatinine clearance < 30 mL/min) • refractory hypokalaemia • hypercalcaemia • concomitant use with ciclosporin and other potent P-gp inhibitors • Hereditary or idiopathic angioedema. **WARNINGS/PRECAUTIONS:** • Avoid use in women planning to become pregnant • Symptomatic hypotension in sodium- and/or volume-depleted patients which should be corrected prior to initiation of therapy. • Caution in patients with serious congestive heart failure and patients with a history of angioedema • Treatment should be discontinued if angioedema or symptoms suggestive of a hypersensitivity reaction/angioedema occur and appropriate therapy and monitoring provided until resolution of signs and symptoms. • Caution is advised when administering Rasilez HCT to patients with renal artery stenosis, renal and liver impairment, renovascular hypertension or systemic lupus erythematosus. • Disturbance of serum electrolyte balance including hypokalaemia, hypochloremic alkalosis, hyponatraemia and hypercalcaemia (monitoring recommended), glucose tolerance and serum levels of cholesterol, triglycerides and uric acid. • Use with caution in patients with aortic and mitral valve stenosis. • Caution with moderate P-gp inhibitors such as ketoconazole. • Caution with concomitant potassium-sparing diuretics, potassium supplements or potassium-containing salts. • Stop treatment in the event of severe and persistent diarrhoea. • Caution in excessive reduction of blood pressure in patients with ischaemic cardiopathy of ischaemic cardiovascular disease. • Caution in driving or operating machinery. • Caution with patients with history of allergy and asthma. • Should not be used during the first trimester of pregnancy • Not recommended in patients below 18 years of age. • Excipients: Contains lactose and wheat starch. **INTERACTIONS:** • Monitoring when used concomitantly with furosemide, lithium, products affected by serum potassium disturbances (eg digitalis glycosides, antiarrhythmics), calcium supplements or calcium sparing medicinal products • Possible interaction with digoxin, St. John's wort, and rifampicin • Meals with high fat content substantially reduce absorption. • Caution when used concomitantly with drugs that may increase potassium levels (eg potassium supplements, heparin sodium) and drugs that decrease potassium levels (eg corticosteroids, ACTH, amphotericin, carbenoxolone, penicillin G, laxatives, salicylic acid derivatives, other kaliuretic diuretics). • Caution if combined with other antihypertensives, curare derivatives, NSAIDs (especially in the elderly), digoxin, antidiabetic agents, allopurinol, amantadine, diazoxide, cytotoxic drugs, anticholinergic agents, cholestyramine and colestipol resins, vitamin D, calcium salts, pressor amines, anitrog medicine, and ciclosporin. • Caution should be exercised on concomitant use with ketoconazole or other moderate P-gp inhibitors (ketoconazole, itraconazole, clarithromycin, erythromycin, amiodarone, leflunomycin). • Grapefruit juice • Alcohol **UNDESIRABLE EFFECTS:** Common: Diarrhoea For the aliskiren component, other reported adverse reactions include: Uncommon: Rash, hyperkalaemia, acute renal failure, renal impairment, oedema peripheral. Rare: Angioedema, haemoglobin decreased, haematocrit decreased, blood creatinine increased. Laboratory values: decrease in haemoglobin and haematocrit, increase in serum potassium. Frequency not known: peripheral oedema. For the hydrochlorothiazide component, other reported adverse reactions include: Aplastic anaemia, bone marrow depression, neutropenia/agranulocytosis, haemolytic anaemia, leucopenia, thrombocytopenia, depression, sleep disturbances, restlessness, light-headedness, vertigo, paraesthesia, dizziness, transient blurred vision, xanthopsia, cardiac arrhythmias, postural hypotension, respiratory distress (including pneumonitis and pulmonary oedema), pancreatitis, anorexia, diarrhoea, constipation, gastric irritation, sialadenitis, loss of appetite, jaundice (intrahepatic cholestatic jaundice), anaphylactic reactions, toxic epidermal necrolysis, necrotising angitis, (vasculitis, cutaneous lupus erythematosus-like reactions, reactivation of cutaneous lupus erythematosus, photosensitivity reactions, rash, arthralgia, weakness, muscle spasm, interstitial nephritis, renal dysfunction, fever. Laboratory values: electrolyte imbalance, including hypokalaemia and hyponatraemia, hyperuricaemia, glycosuria, hyperglycaemia, increases in cholesterol and triglycerides **LEGAL CATEGORY:** POM **PACK SIZES:** 7, 28 film-coated tablets **MARKETING AUTHORISATION HOLDER:** Novartis Europharm Limited, Wimblehurst Road, Horsham, West Sussex, RH12 5AB, United Kingdom. **MARKETING AUTHORISATION NUMBERS:** Rasilez HCT 300/12.5 mg - EU/1/08/491/041-060. Rasilez HCT 300/25 mg - EU/1/08/491/061-080. Please refer to Summary of Product Characteristics (SmPC) before prescribing. Full prescribing information is available on request from Novartis Pharma Services Inc, Representative Office Malta, P.O. Box 124, Valletta, VLT 1000, Malta. Tel +356 22963217. (2011-MT-02-RASILEZ HCT Feb 2011)

 **NOVARTIS**

References: 1. Palatini P, Jung W, Shlyakht E, et al. *J Hum Hypertens* 2010; 24:93-103; published online 21 May 2009. 2. Villamil A, Chrysant SG, Calthun D, et al. *J Hypertens* 2007; 25:217-226

ALBERT CILIA-VINCENTI

THE SERIES

# Healing & Disease Reversal

This series reviews Dean Ornish's evidence-based claims of healing & disease reversal by dietary and lifestyle changes. He is a California University Professor of Medicine in San Francisco. This instalment discusses "bad" fats.

**T**rans-fatty acids, partially hydrogenated fats and saturated fats ("bad fats") should be avoided. Many food manufacturers continue to use them because they increase their products' shelf-life, even though they may decrease the "shelf-life" of the people who consume them.

Saturated fats are found mainly in animal fats – in meats (including chicken and other poultry), full-fat dairy produce (milk, cream, cheese and ice-cream), egg yolk and (to a lesser extent) seafood. Some plant foods, such as coconut and palm oils, are also rich in saturated fats.

When vegetable oils are heated in the presence of hydrogen, both partially hydrogenated and trans-fatty acids are produced. Partially hydrogenated fats are as disease-promoting as saturated fats.

Trans-fatty acids are especially bad for health. They are found predominantly in commercially prepared baked foods, margarines,

snacks, fast foods, processed and fried foods. Trans-fatty acids are probably worse for cholesterol levels than saturated fats because they raise bad LDL and lower good HDL. It's wise to avoid both saturated fats and trans-fatty acids.

Besides raising total and LDL cholesterol levels, saturated, partially hydrogenated fats and trans-fatty acids promote inflammation and are strongly linked with increased risk of coronary heart disease, stroke, diabetes, many types of cancer, and other chronic diseases.


A recent scientific presentation<sup>1</sup> also claimed that diets rich in trans-fatty acids may cause a redistribution of fatty tissue to the abdomen (the worst place to store fat for both health and appearance) and lead to a higher body weight, even when total calorie intake is the same.

Processed food labels now list both saturated and trans-fatty acids. Also, the more hydrogenated an oil is, the


harder it will be at room temperature. Soft tub margarines usually have less hydrogenated fats and trans-fatty acids than harder margarines that come in bars.

Professor Ornish began consulting with McDonald's in 1999, to help them develop more healthy foods. His first recommendation was to remove trans-fatty acids from their products. This was quite a challenge for McDonald's to preserve the flavour and texture of its French fries, but they found a way to do it. He then helped them develop a Fruit & Walnut Salad, containing apple, grapes, walnuts and low-fat yogurt – McDonald's is now the largest purchaser of apples in the world. He also helped them develop an Asian Salad, containing sixteen types of greens, soybeans, almonds, mandarin oranges, peas and red bell peppers.

He has also helped PepsiCo and Safeway Supermarkets, and is pleased that when making more healthy foods becomes good business, it becomes sustainable. At PepsiCo, for example, more than two thirds of revenue growth now comes from its good-for-you and better-for-you products.

It takes time for companies of this size to change their product range to include more healthy foods, but he feels it's happening more quickly than he expected, and is proud of the impact it's having worldwide in making it more convenient and fun to eat more healthily. 

Professor Ornish began consulting with McDonald's in 1999, to help them develop more healthy foods. His first recommendation was to remove trans-fatty acids from their products



## Reference

1. Rudel LL, Kavanagh K. Tran fat leads to weight gain even on same total calories. The 66<sup>th</sup> Annual Scientific Sessions of the American Diabetes Association, Washington, DC, 2006.

# THE POWER OF 3

## PROVEN AGENTS IN A SINGLE PILL

For Patients who need **MORE** than dual therapy to reach Blood Pressure goal<sup>1</sup>



Single-pill power superior to dual therapy<sup>2,3</sup>



Single-pill power sustained for 24 hours<sup>4</sup>



Single-pill power with a well established safety profile<sup>2</sup>

References: 1. Mancia G, Laurent S, et al. *Blood Pressure* 2009;18:308-347. 2. Calhoun DA, Lacourcière Y, Chiang YT, Glazer RD. *Hypertension* 2009;54:32-39. 3. Lacourcière Y, Glazer R, Crikelair N, Yen J, Calhoun D. Poster presented at: 19th Scientific Meeting of the ESH; 12-16 June 2009; Milan, Italy. 4. Lacourcière Y, Glazer R, Yen J, Calhoun D. Poster presented at: 19th Scientific Meeting of the ESH; 12-16 June 2009; Milan, Italy.

**EXFORGE HCT**  
amlodipine besylate/valsartan/hydrochlorothiazide

GREAT DROPS JUST GOT BETTER

NOVARTIS

Exforge® HCT (amlodipine besylate /valsartan/hydrochlorothiazide) 10/160/25mg and 10/320/25mg Film-coated tablets

**PRESENTATION:** Film-coated tablets containing: 10 mg amlodipine as amlodipine besylate, 160 mg valsartan and 25 mg hydrochlorothiazide or 10 mg amlodipine as amlodipine besylate, 320 mg valsartan and 25 mg hydrochlorothiazide. **INDICATION:** Treatment of essential hypertension as substitution therapy in adult patients whose blood pressure is adequately controlled on the combination of amlodipine, valsartan and hydrochlorothiazide (HCT) taken either as three single-component formulations or as a dual-component and a single-component formulation. **DOSAGE:** One tablet of Exforge HCT 10/160/25 mg or 10/320/25 mg daily. **CONTRAINDICATIONS:** ♦ Known hypersensitivity to the active substances, to other sulfonamides, to dihydropyridine derivatives, or to any of the excipients ♦ Second and third trimesters of pregnancy ♦ Hepatic impairment, biliary cirrhosis or cholestasis ♦ Severe renal impairment (creatinine clearance < 30 mL/min) ♦ Anuria ♦ patients undergoing dialysis ♦ Refractory hypokalaemia ♦ Hyponatraemia ♦ Hypercalcaemia ♦ Symptomatic hyperuricaemia. **WARNINGS/PRECAUTIONS:** ♦ Risk of hypotension in sodium- and/or volume-depleted patients (correction is recommended prior to administration of Exforge HCT). ♦ Caution is advised when administering Exforge HCT to patients with renal impairment or systemic lupus erythematosus. ♦ No data available in patients with unilateral or bilateral renal artery stenosis, stenosis to a solitary kidney or after recent kidney transplantation ♦ Disturbance of serum electrolyte balance (monitoring recommended), glucose tolerance and serum levels of cholesterol, triglycerides and uric acid. ♦ Not recommended in patients below 18 years of age and in patients with primary hyperaldosteronism ♦ Beta-blocker withdrawal should be gradual. ♦ Caution in elderly and in patients with hepatic impairment or biliary obstructive disorders. ♦ Caution in patients with heart failure and coronary artery disease ♦ As with all other vasodilators, special caution in patients suffering from aortic or mitral stenosis, or obstructive hypertrophic cardiomyopathy. ♦ If a photosensitivity reaction occurs during treatment, it is recommended to stop the treatment. ♦ Not recommended during the first trimester of pregnancy. Avoid use in women planning to become pregnant and while breast-feeding. ♦ Caution when driving or using machinery. **INTERACTIONS:** ♦ Monitoring recommended when used concomitantly with lithium. ♦ Caution when used concomitantly with drugs that may increase potassium levels. ♦ Caution if combined with other antihypertensives, curare derivatives, NSAIDs, corticosteroids, ACTH, amphotericin, carbenoxolone, Penicillin G, salicylic acid derivatives, digoxin, CYP3A4 inhibitors and inducers, antidiabetic agents, allopurinol, probenecid, sulfipyrazone, pressor amines, amantadine, diazoxide, cytotoxic drugs, anticholinergic agents, methyldopa, cholestyramine, cholestipol resins, vitamin D, calcium salts, carbamazepine and ciclosporin, alcohol, anaesthetics and sedatives. **ADVERSE REACTIONS:** ♦ Exforge HCT (amlodipine/valsartan/HCT): Common: hypokalaemia, headache, dizziness, hypotension, dyspepsia, pollakiuria, oedema, fatigue. Uncommon: anorexia, hypercalcaemia, hyperlipidaemia, hyperuricaemia, hyponatraemia, insomnia/sleep disturbances, abnormal coordination, postural and exertional dizziness, dysgeusia, lethargy, paraesthesia, peripheral neuropathy, neuropathy, somnolence, syncope, visual disturbance, vertigo, tachycardia, orthostatic hypotension, phlebitis, thrombophlebitis, cough, dyspnoea, throat irritation, abdominal discomfort, upper abdominal pain, breath odour, diarrhoea, dry mouth, nausea, vomiting, hyperhidrosis, pruritus, back pain, joint swelling, muscle spasm, muscular weakness, myalgia, pain in extremity, elevation of serum creatinine, acute renal failure, erectile dysfunction, abasia, gait disturbance, asthenia, discomfort, malaise, non cardiac chest pain, increased blood urea nitrogen, increased blood uric acid, decreased serum potassium, weight increase. ♦ Additional adverse reactions with amlodipine monotherapy: Common: palpitations, flushing. Uncommon: mood swings, tremor, tinnitus, rhinitis, change of bowel habit, alopecia, exanthema, purpura, skin discoloration, arthralgia, micturition disorder, nocturia, gynaecomastia, pain, weight decrease. ♦ Additional adverse reactions with HCT monotherapy: Common: increased lipids. Uncommon: hypomagnesaemia, decreased appetite, urticaria. Rare: thrombocytopenia, hyperglycaemia, depression, arrhythmia (including bradycardia, ventricular tachycardia and atrial fibrillation), constipation, intrahepatic cholestasis, jaundice, photosensitivity reaction, renal failure and impairment, glycosuria. **LEGAL CATEGORY:** POM **PACK SIZES:** Packs of 28 film-coated tablets **MARKETING AUTHORISATION HOLDER:** Novartis Europharm Limited, Wimblehurst Road, Horsham, West Sussex, RH12 5AB, United Kingdom. **MARKETING AUTHORISATION NUMBER:** Exforge HCT 10 mg/160 mg/25 mg - EU/1/09/569/038 Exforge HCT 10 mg/320 mg/25 mg - EU/1/09/569/050 Please refer to Summary of Product Characteristics (SmPC) before prescribing. Full prescribing information is available on request from Novartis Pharma Services Inc., Representative Office Malta, P.O. Box 124, Valletta, VLT 1000, Malta. Tel: +356 22983217 2010-MT-01-EXFH-16-OCT-2009

Check out the Stresam eQuiz on [www.thesynapse.net](http://www.thesynapse.net).

We encourage all doctors to participate. The winning prize is a meal for two persons at the exclusive Grill 3301 St Julians.

## Quest eQuiz winner



Quest Vitamins in collaboration with Pharma.MT Ltd would like to congratulate **DR JOHN XUEREB CURMI** on winning the Quest Vitamins E-Quiz. He has been presented with a 6 month free supply of Super Once a Day Timed Release. This supplement is an all in one high potency Multivitamin formula which provides 17 vitamin and vitamin-related ingredients, 12 minerals and 12 additional factors in a 6 hour timed release preparation. Dr Xuereb Curmi has also been presented with a Quest Excellence IQ Manual and Quest Product Guide, which both contain valuable information for all health care professionals who advise and prescribe vitamins, minerals, herbs and other health supplements.

**Locum pharmacist** required to work alternate Saturdays. Please contact Mr. Malcolm Muscat on 7959 5398.

Krypton Chemists Ltd., St. Andrews, require a **locum pharmacist** part time. The ideal candidate should have an outgoing personality and be able to work on their own initiative. The pharmacist will be part of a team with two other pharmacists, so flexibility is available. An attractive remuneration will be offered to the right applicant. Interested applicants are required to send their CV to [imports@kryptonchemists.com](mailto:imports@kryptonchemists.com) and to contact the Managing Director on 2137 8888 or 2137 0141.

**Pharmacist wanted** to work in community pharmacy full time or mornings only. Please contact managing pharmacist on telephone: 21662270 / 99452507.

### Equipment for sale

- 2 Visiting Couches - Black with head rests @ €200 each
- ECG Machine - (NIHON KOHDEN, 12 lead 3 Channel) @ €700
- EEG Machine - (NIHON KOHDEN) @ €7000

Items mentioned above are in perfect condition, and have been bought some 5 years ago.

Contact [ed\\_apap@gmail.com](mailto:ed_apap@gmail.com).

### Clinic available

Clinic available for GP / Specialist mornings & evenings Monday to Saturday in well-established pharmacy in St. Joseph High Road, Hamrun.

Please contact Ms Sciberras on 79498337 / 21227682.

### Pharmacist Vacancy

Pharmacist wanted to work in community pharmacy full-time or mornings only. Please contact managing pharmacist on 21662270 / 99452507.

# Lyprinol®

**Move Without Pain**

*The All Natural  
and Safe Solution*

*No Risks! No Side Effects!  
Only Positive Benefits*

- Increased Heart Health & Joint Mobility
- Relieve the pain of arthritis & Joints
- Reduce Joint Inflammation & Swelling
- Better Breathing
- Potentially increased Brain Function
- Reduces Asthmatic Symptoms

• 150 times more effective than fish oil • No repeats  
 • Very easy to swallow • Suitable for diabetics  
 • Suitable for children over 3 years of age  
 • Lyprinol is free of Gluten, Lactose, Sugar and Preservatives




Trade Enquiries:  
**a.t.g. co. ltd.**  
*dedicated to well being since 1992*

Tel: +356 2124 2017 | E-mail: [ales@atg.com.mt](mailto:ales@atg.com.mt)  
[www.atg.com.mt](http://www.atg.com.mt)

Available in all leading pharmacies in Malta and Gozo

## Valsotens eQuiz Winner

The overall prize winner for  
The Synapse Valsotens eQuiz was  
**DR MARK ANTHONY VASSALLO**



Valsotens contains valsartan which is an angiotensin II antagonist. It lowers blood pressure by providing specific and selective blockade of action of Angiotensin II at the AT<sub>1</sub> receptor.

The specificity of action of Angiotensin II antagonists avoids some of the unwanted effects of other drugs that inhibit the Renin-Angiotensin-System, such as interruption of bradykinin-onward metabolism leading to cough with ACE inhibitors. Valsotens is available in 2 doses: 80mg and 160mg. Each dose is available in blister packs of 28 film-coated tablets.

Valsotens is indicated for:

- Hypertension
- Recent-myocardial infarction: improves survival following recent (12 hours – 10 days) myocardial infarction in clinically stable patients with signs, symptoms or radiological evidence of left ventricular failure and/or with left ventricular systolic dysfunction.

### Errata Corrige 2/11

The article by Prof Albert Cilia-Vincenti should have included the following reference: Ebbeling CB, Leidig M, Feldman H, Lovesky M and Ludwig D. Effects of a low-glycemic load vs low-fat diet in obese young adults.

*J Am Med Ass* 2007; 297 (19): 2092.

The error is regretted.

Think of it as your own automatic personalised recall system

You and your patients only stand to gain!




Contact us now  
and we will help  
you improve  
healthcare  
delivery to your  
patients...

Tel: 21453973  
Email: [helpdesk@sms4health.com](mailto:helpdesk@sms4health.com)  
Website: [www.sms4health.com](http://www.sms4health.com)



*Avoid clogs*



## **Agregex**

*Clopidogrel  
75mg film-coated tablets  
Antithrombotic Agent*

*For further information please refer to the Full Summary of Product Characteristics  
or to our website: [www.actavis.com.mt](http://www.actavis.com.mt)*

 **actavis**  
creating value in pharmaceuticals

# Management of consumers' pharmaceutical waste in a pharmacy setting - Part I

The management of pharmaceutical waste is increasingly becoming of great concern to our society since no official program currently exists. This study proposes a management plan for consumer's pharmaceutical waste in a pharmacy setting in Malta. It involves disposing of unwanted solid dosage-forms in their original packaging in waste containers which will be installed in front of pharmacies, and returning hazardous medicines and other dosage-forms directly to the pharmacist to reduce hazards from mixing incompatible products.

## Key words

Pharmaceutical waste management; pharmacy; consumers; waste containers.

## Introduction

Ever since joining the European Union, Malta has been trying extensively to conform to European standards as requested by legislation. Improvements have been registered in several sectors; however the research in the management of waste pharmaceuticals has so far remained lacking.<sup>1</sup>

## Aims

This report draws findings of a scientific study that has been undertaken by the authors and aims to provide a programme and a guidance proposal plan for the management and disposal of pharmaceutical waste produced by consumers. The process of designing and implementing a successful pharmaceutical waste management program, being highly

interdisciplinary, has to be acceptable for pharmacies, competent authorities and consumers alike.<sup>2</sup>

## Educational programs

Educational programs should be set up to educate consumers on the threats resulting from the irresponsible disposal of pharmaceutical waste. The proposed programs should educate users on new, well thought-out and science-founded procedures.<sup>3</sup> The benefits should be highlighted to encourage people to make an effort to abide by these guidelines.

Such educational programs include:

- Promoting the campaign on television programs including waste management 'spots' on TV;
- Advertisements on published media;
- Distribution of leaflets from community pharmacies;
- Other advertising materials including banners and posters.


## Consumers' pharmaceutical waste

Examples of consumers' pharmaceutical waste include:

- Expired medicines in households;
- Medicines discontinued by manufacturers due to a negative risk-benefit ratio;
- Medicines not consumed by patients;
- Medicines discontinued because the doctor prescribed new medication;
- Medicines of deceased patients.

Since pharmaceutical waste production in nursing homes is significantly higher than in households, this scheme is not applicable. In this setting, clinical waste is also produced and thus nursing homes should contract a private waste management company to manage their pharmaceutical and clinical waste. Hospitals have their in-house waste management procedures.

Figure 1: Open Pharmaceutical Waste Container



Household waste medicines are found in various dosage forms; the most common being tablets, capsules, caplets, syrups, suspensions, inhalers, suppositories, pessaries, creams, ointments, sprays, drops, powder for oral constitution and patches.<sup>4</sup>

Syringes are not included in this scheme. Yellow UN approved containers should be distributed to frequent syringe users from healthcare centres. When the yellow bin is full, consumers should return the bin to the centre and they are given an empty one. The bins are then collected from the healthcare centres by waste carriers.

### Hazardous waste

Section 4(2) of the Waste Management Act of 2001 defines Hazardous Waste as having certain properties which can be harmful to human health and the environment. "These properties are corrosive, explosive, toxic, flammable, oxidizing, irritant, harmful, carcinogenic, infectious, teratogenic, mutagenic and ecotoxic."<sup>5</sup>

Medicines may have a range of hazardous properties, but only those with one of the hazardous properties denoted by "H codes" will be classified as a hazardous waste.<sup>5</sup> Other medicines are not hazardous within the meaning of the legislation, but they are hazardous if released in the environment.

### Hazardous medicines

The European Waste Catalogue<sup>5</sup> lists wastes from human healthcare in Chapter 18. Medicines consisting of or containing dangerous substances together with cytotoxic and cytostatic medicines are considered to be hazardous. Hazardous medicines include chemotherapeutic agents, antivirals, hormones and some bioengineered medicines.<sup>6</sup> Most of the medicines available in community pharmacies are fairly inert and pose no particular danger to staff or waste carriers who collect the waste. This is because most medicines are individually packed tablets or capsules. The packaging provides

**Table 1:** Hazardous medicines available in pharmacies (Source: Malta Medicines List)  
This table is only a summary of the full version which is part of the scientific study entitled 'Waste Management in Pharmacy'

Anti-acne preparations		
Active ingredient	Trade Name	Formulation
Clindamycin 10mg/ml	Dalacin T	Topical solution
Isotretinoin 10mg, 20mg	Decutan	Capsule, Soft
Anti-bacterials for systemic use		
Clavulanic Acid 125mg; Amoxicillin 500mg	Augmentin-Duo 500mg/125mg	Film-Coated Tablet
Cefaclor 500mg	Ceclor	MR tablets
Antibiotics and chemotherapeutics for dermatological use		
Retampulin	Altargo	Ointment
Mupirocin 2% w/w	Bactroban	Topical ointment
Intestinal anti-inflammatory and anti-infective agents		
Mesalazine 400mg	Pentacol	Gastro Resistant Tablet
Sulfasalazine 500mg	Salazopyrin	Suppositories
Anti-gout preparation		
Allopurinol 100mg	Allopurinol	Tablet
Colchicine 500mcg	Colchicine	Tablet
Anti-neoplastic agents		
Methotrexate 2.5mg	Methotrexate	Film-Coated Tablet
Imatinib 100mg	Imatinib	Tablets
Endocrine therapy		
Letrozole 2.5mg	Femara	Tablet
Tamoxifen 20mg	Nolvadex D	Film-Coated Tablet
Ophthalmologicals		
Tobramycin 3mg/ml	Tobrex	Eye Drops, Solution
Aciclovir 3% w/w	Zovirax	Eye Ointment
Sex hormones and modulators of the genital system		
Estradiol 25mcg	Vagifem	Vaginal Tablet
Ethinylestradiol 0.03mg, Drospirenone 3mg	Yasmin	Film-Coated Tablet
Thyroid therapy		
Levothyroxine Sodium 100mcg	Eltroxin	Tablet
Carbimazole 5mg	NeoMercazole 5	Tablet

satisfactory protection for anyone handling the container. However if the solid formulation is altered by, for example, being crushed, they could pose a risk. Blister packaging also protects against mixing incompatible products and they do not pose a risk to the environment provided that they are incinerated properly.<sup>2</sup>

Pharmacies should develop a list of hazardous medicines that are found in the pharmacy, including medicines

available from the Pharmacy of Your Choice (POYC Scheme) so as to be aware of any risks involved during collection.

Table 1 is a short template of a list of medicines available in pharmacies which are considered to be hazardous, that was compiled as part of the scientific study undertaken. This list should be modified according to what is available at each respective pharmacy. [cont in issue 4/11](#)

# Becoming a dentist in the 50s

She was not quite convinced she really wanted to do the interview. "I've been through the story so many times... isn't it a bit boring?" I admitted I had never heard it before, so at least, for my sake, it was worth re-telling once again. Carmen Joslin, very pragmatically tells me she was the first female dentist in Malta and shrugs it off as nothing special. After I hear her story, I'm not so convinced.

Mrs Joslin or rather Dr Joslin nee Attard, graduated in 1960 but her introduction to the possibility of studying dentistry actually began way before that. As a child she was born in in the then simple quiet village of Balzan, just a few years before World War II broke out. "I was born in 1938 and my sister and I were sent to school in Valletta to St Joseph High School. My parents were not wealthy but they both instilled in us the early encouraging notion of going to University. You have to understand that back then, University was not free as it is today and it was expensive for that time. Moreover, nobody really encouraged daughters to go to University, as it was only the boys who were considered in that regard. It was a new thing to have a girl go for a university education and in that sense I still marvel at how avant-garde my parents were in their thinking."

Another unique habit for the times was that as part of her parents'

resolve to keep the daughters' teeth as healthy as possible, annual check-ups at the dentist were the norm. It was during one of these routine visits that Carmen Joslin became fascinated with the idea of dentistry. "I was sitting on the dentist's chair with my mouth wide open, when my father flippantly mentioned that we (my sister and myself) wanted to go to university and asked for the dentist's advice in this regard. It was his and my mother's conviction in the first place, but anyway, that is beside the point. As part of his reply, the late Dr Tony Demajo who was our dentist at the time commented that there were no female dentists and this comment remained ingrained in my head. One of my characteristics as a person is that I don't want to be herded with the rest – I prefer doing my own thing, and the notion of being a first or at least one of the first, enticed me. From that day onwards, the thought never left my mind. There were some hurdles I had to over come. An 'O' level pass in Matriculation Maltese was a requirement for entry to university. Maltese was never taught at my school and so I had to study it privately after school hours."

Even whilst the mention of university study was frequent, Carmen Joslin admits she did not really know what university was all about, didn't even know where the building proper was situated. But by the time she was



through her 'O'Levels and ready to start attending the university which was in Valletta at the time, she found she had to first attend a two year course in Science and Art, topics not taught by the nuns. The two years meant to give her and others grounding in the basic notions of the sciences and other subjects. It was a day and age when youngsters had no real knowledge of science, unlike today when most know at least the basic way their body functions and something about the organs it includes. As Dr Joslin succinctly puts it, "We didn't even know we had ovaries, let alone the rest of it."

This was in 1954 when the total intake at university for the year was very small and very few of these were females. It was considered a waste of time and money to educate females to that educational level as they would be getting married in a few years time anyway.

"The first two years I spent at University were very different to my school days. We girls felt like birds that had been given their freedom. When the two years were over I applied for the four year course leading to a degree in Dental surgery, the only



...married women were not accepted as employees of [St Luke's] hospital

# Medicool, Advanced Medicinal Transportation

Keep-Cool  
Carry Cases

Micro-Refrigerators  
For Car Use  
12V/240V

Ideal for both the Patient and  
the Medical Professional

Take your medicines with you, wherever you go!  
**NOW YOU CAN!**

**JAMESCO**  
TRADING COMPANY LIMITED  
Tel: 21314333

Call or Email for more Info:

customer@jamescotrading.com  
www.jamescotrading.com  
21314333

woman who did so. I must admit that throughout all of my studies, I never found one single professor or anybody else for that matter, who discouraged me. My teachers were Professor Edigio Lapira, and Professor Joseph Mangion, besides many others.”

The first two years were spent studying anatomy, physiology, bacteriology amongst other subjects. The last two years were intensive. Dental surgery, pathology .and conservative dentistry were two subjects and they also studied general medicine and general surgery apart from others. Carmen Joslin still marvels at how people believe a dentist is just somebody who pulls out teeth. It was an exciting time for her – and at the end of it, only seven students had passed in her course.

She placed third in the course and remained the only lady on board, sharing graduation with other dental surgeons – Dr Herbert Messina Ferrante, Dr Victor Demicoli, Dr George Otibah (Nigerian), Dr Joe Bartolo, Dr Ronald Delia and Dr (now Professor) George Zarb.

In the meantime, she had met the young man who would become her husband – Sidney Joslin. “We met



Post graduation photo who graduated in 1960. From left to right - Drs Herbert Messina Ferrante, Victor Demicoli, George Otibah (Nigerian), Carmen Joslin, Joe Bartolo, Ronald Delia. Professor George Zarb was also part of my group but had already left for Canada to start his post-graduate studies

a major let-down for her. Her first employment was within St Luke’s Hospital where she spent a relatively short period of time. She formed part of a team and had the benefit of being able to consult with others where difficult cases were concerned. Whilst that lasted it was good. But once word transpired that she planned to get married, she was asked to present herself to administration. “They

teamwork experienced previously. Once the babies were born, she started working on reduced hours, drastically diminishing the time spent at the dental clinic. “With three babies on board, I opted out. Babies and small children require a mother’s attention continuously. Eventually when the policies at St Luke’s Hospital changed, I was invited to resume my work there, but by that time quite some years had elapsed and I did not feel up to it. Mostly because I felt I was not up to date with the latest techniques, and dentistry being the practical profession that it is, you cannot rely on theory only.”

Today Dr Joslin is mother of four adult siblings and grandmother of five and keeps busy with her garden, reading and spending time with animals which are another love of her life. She has also discovered the thrills of computing and admits she spends many interesting hours engrossed in the effort. “Those who know me as Dr Joslin still refer to me that way but most people know me simply as Dr Joslin and that suits me just fine. It seems a long time ago but what I trained for has certainly stood me in good stead – the formation you get at university is important. I would have probably kept my job at the hospital had I been allowed to do so. But that was the way things worked out back then...” S

## Her first dismay was to find that women received only three quarters of what their male counterparts received in terms of salary

at university and had been seeing each other for some four years when I graduated. It was not done at the time. People who became engaged, were married off soon afterwards. But Sidney went to the UK to study and we initially planned that we would marry and I would join him and continue working as a dentist there. It never materialised as he came back to Malta and we settled on the island. But in the meantime, once I finished university which had been one of the most enjoyable periods of my life... problems started in earnest.”

Her first dismay was to find that women received only three quarters of what their male counterparts received in terms of salary. This was already

politely but matter-of-factly suggested I present a letter of resignation, telling me that married women were not accepted as employees of the hospital. I was appalled. It was news to me and I refused to resign, telling them that if they didn’t want me there anymore, it was up to them to fire me. They eventually did so by means of a letter diplomatically telling me my services were no longer required. I was dismayed and utterly appalled by the discrimination of it all.”

Dr Joslin did not give up but went into private practise with the very same dental surgeon who had seen her teeth as a child – Dr Demajo. However she never took to the private practise scenario much, missing the

# Idéos®

Calcium/Vitamin D<sub>3</sub> 500mg/400IU

## A new Idea of Osteoporosis



**2 tablets per day**

**Chewable tablets  
Lemon flavour**

**Convenient for patients with:**  
- Diabetes  
- High blood pressure

**Can be associated with any specific  
treatment of osteoporosis**

**IDEOS®**, chewable tablets. **COMPOSITION\***: Elemental calcium 500 mg (corresponding to 1250 mg of calcium carbonate), cholecalciferol (Vitamin D<sub>3</sub>) 400 IU (corresponding to 4 mg of cholecalciferol concentrate, powder form). **PHARMACEUTICAL FORM** : Chewable tablets. **THERAPEUTIC INDICATIONS** : Vitamin D and calcium deficiency correction in the elderly, Vitamin D and calcium supplementation, as an adjunct to specific therapy for osteoporosis, in deficient patients or in patients with a high risk of vitamin D and calcium deficiency. **POSOLGY AND METHOD OF ADMINISTRATION** : Oral use. For adults only. Suck or chew the tablets. One tablet twice a day. **CONTRAINDICATIONS**: Hypersensitivity to one of the constituents. Hypercalcaemia, hypercalciuria, calcium lithiasis. In patients where prolonged immobilisation is accompanied by hypercalcaemia and/or hypercalciuria, vitamin D and calcium treatment should only be resumed when the patient becomes mobile. **SPECIAL WARNINGS AND PRECAUTIONS FOR USE** : In case of long term treatment, the urinary calcium excretion must be monitored and treatment must be reduced or temporarily suspended if urinary calcium excretion exceeds 7,5 mmol/24 h, i.e. 300 mg/24 h. In case of simultaneous treatment with digitalis glycosides, diphosphonates, sodium fluoride, thiazide diuretics, tetracyclines, see "Interactions". Take into account the dose of vitamin D per tablet (400 IU) in case of simultaneous treatment with another vitamin D preparation. Administration of supplementary vitamin D or calcium should be done under medical surveillance with monitoring of calcaemia and calciuria. The product should be prescribed carefully to patients suffering from sarcoidosis because of the potential increase of vitamin D metabolism into its active form. For these patients calcaemia and calciuria should be monitored. In case of renal insufficiency, adapt the dosage according to creatinine clearance. **INTERACTION WITH OTHER MEDICAMENTS AND OTHER FORMS OF INTERACTION\***: Digitalis glycosides, thiazide diuretics, oral tetracyclines, - diphosphonates, sodium fluoride, high doses of vitamin D, phenytoin, barbiturics, glucocorticoids. **PREGNANCY\***: Ideos can be used during pregnancy and lactation. However, the daily dose should not exceed 1500 mg of calcium and 600 IU of vitamin D<sub>3</sub>. During pregnancy, overdose of cholecalciferol and vitamin D should be avoided. **LACTATION**: Vitamin D and its metabolites enter the breast milk. **UNDESIRABLE EFFECTS** : Hypercalciuria and exceptionally hypercalcaemia, in case of long term treatment in high dosage: constipation, flatulence, epigastric pains, diarrhoea. **OVERDOSE\***. **PHARMACOLOGICAL PROPERTIES\*** : VITAMIN D-CALCIUM SUPPLEMENT (Drug acting on calcium balance - Alimentary tract and metabolism). **PHARMACEUTICAL FORMS AND PRESENTATIONS**: . Box of 60 tablets in tube of 15 tablets. MA number: MA093/00201 **MARKETING AUTHORISATION HOLDER**: LABORATOIRE INNOTECH INTERNATIONAL 22 Avenue Aristide Briand 94110 ARCUEIL-FRANCE. TEXT REVISION DATE:..14 April 2010. \*For complete information, please refer to the therapeutic references.

# BIAFINE®

## EMULSION

Johnson & Johnson



SECONDARY ERYTHEMA DUE TO  
RADIOTHERAPY TREATMENTS



FIRST & SECOND DEGREE BURNS  
INCLUDING SUNBURNS



ALL NON-INFECTED  
CUTANEOUS WOUNDS

including:

- ✓ Stage I - IV Pressure Ulcers
- ✓ Diabetic ulcers
- ✓ Venous & Ischaemic ulcers
- ✓ Hypertensive Ulcers
- ✓ Chronic & Traumatic wounds.




Tube of 93g

ENQUIRIES: To request samples  
or more information about this product,  
kindly contact Medical Representative on  
Tel. 27010276 / 9984 5587  
E-mail: kemifsud@go.net.mt

**TRADE NAME:** BIAFINE®, cutaneous emulsion **QUALITATIVE AND QUANTITATIVE COMPOSITION:** TROLAMINE 0.670 g; **PHARMACEUTICAL FORM:** Cutaneous Emulsion. **CLINICAL PARTICULARS:** Therapeutic indications: Erythema secondary to radiotherapy. First- and second-degree burns and any other uninfected skin wounds. **Dosage and method of administration:** - Erythema secondary to radiotherapy: two to three applications daily, at regular intervals, ensuring penetration by gentle massage. - Second-degree burns and other skin wounds: after cleaning the wound, apply a thick layer, to the lesion and around its periphery, and renew applications in order to permanently maintain excess emulsion on the lesion. If required, cover with a moist compress and complete the dressing. Do not use dry absorbent dressings. - First-degree burns: apply a thick coat until no further emulsion can be applied. Massage gently to ensure penetration. Repeat two to four times daily. **Contraindications:** - known allergy to one of the constituents of the preparation - hemorrhagic wound, - infected lesion. **Special warnings and special precautions for use:** Special warnings: In the event of a second-degree burn or uninfected skin wound, management depends on the extent of the lesion, its location, the patient's age and history, the associated lesions and the etiology. **Special precautions for use:** This medicinal product is not a sun screen. **Interactions with other medicinal products and other forms of interaction:** None known. **Pregnancy and lactation:** If you are pregnant or breast feeding, you should always seek the advice of your doctor or pharmacist before using any medication. **Effects on the ability to drive and use machines:** Not Applicable. **Adverse effects:** Sometimes, moderate and transient pain, of the stinging type, after application. Rarely, contact allergy. **Overdose:** Not Applicable. **PHARMACOLOGICAL PROPERTIES:** Pharmacodynamic properties: SKIN PROTECTIVE (D: Dermatology). **Pharmacokinetic properties:** Not applicable. **Preclinical safety data:** Not applicable. **PHARMACEUTICAL PARTICULARS:** List of Excipients: Ethylene glycol monostearate 5.450g; Stearic acid 3.625 g; Cetyl palmitate 0.350g; Solid paraffin 1.600 g; Liquid paraffin 6.850 g; Perhydroqualene 1.500 g; Propylene glycol 2.300 g; Avocado oil 1.000 g; Trolamine and sodium alginate 0.702 g; Potassium sorbate (quantity expressed as sorbic acid) 0.100 g; Sodium methyl parahydroxybenzoate (E 219) 0.100g; Sodium propyl parahydroxybenzoate (E 217) 0.050g; Yerbatoone flavoring\* 0.130g; purified water q.s. 100 g, per 100 g of emulsion for cutaneous application \* *Composition of the Yerbatoone flavoring:* essential oil of Brazil orange, essential oil of galbanum, essential oil of petitgrain, essential oil of lemongrass, 4-acetyl-6-(1,1-dimethylethyl)-1,1-dimethylindane, para-tert-butylcyclohexyl acetate, 4-(4-hydroxy-4-methylpentyl)-3-cyclohexene-10-carboxaldehyde, natural citral, limonene, benzyl benzoate, phenylethyl alcohol, musk ketone, phenylethyl acetate, methylionone, dipropylene glycol. **Incompatibility:** Not applicable **Shelf life:** 3 years. **Special precautions for storage:** Do not store at a temperature less than 0°C. **Nature and contents of container:** Aluminium tube containing 9, 93 or 186 g. The tube has an interior epoxyphenolic coating and is closed by a high-density polyethylene cap. **Instructions for use, handling:** No specific requirement **MARKETING AUTHORIZATION HOLDER:** JOHNSON & JOHNSON CONSUMER FRANCE S.A.S. 1, rue camille Desmoulins, 92130 Issy-Les-Moulineaux, France **PRESENTATION AND ADMINISTRATIVE IDENTIFICATION NUMBER:** MA065/00101 **DATE OF APPROVAL/RENEWAL:** 2nd March 2007 **DATE OF REVISION:** 1st August 2010



# What is Doppler ultrasound?



**Figure 1:** Normal common carotid arterial Doppler ultrasound showing a colour Doppler image (a) and a spectral Doppler depiction (b).

The term *Doppler* should be capitalized because it refers to Christian Johann Doppler, an Austrian physicist (1803–1853). He described a phenomenon whereby the frequency of sound changes when it is reflected off a moving object. If the object is moving towards the observer, reflected sound frequency is increased (blue shift), while if the reflecting object is moving away from the observer, the sound frequency decreases (red shift). This is analogous to a moving ambulance with its siren on; the siren pitch increases as ambulance approaches and decreases as it recedes.

The Doppler Effect may be used also in ultrasound. With normal grey scale ultrasound we rely on amplitude of reflected sound waves and the


reflectivity of anatomic structures is proportional to the intensity (or amplitude) of the reflected sound and hence brightness on the scanner display. If frequency (rather than amplitude) is analysed, one can detect motion on the basis of Doppler shifts. These frequency shifts can be quantified reasonably accurately according to direction, velocity and also acceleration, all of which provide specific signatures that help identify both normal and diseased blood vessels.

Frequency shifts in ultrasound may be displayed in two fashions; they may be presented on a graphical plot showing velocity on the y axis and time on the x axis, this is called spectral Doppler ultrasound, or they may be colour coded with blue or red


indicating direction of flow towards or away from the ultrasound probe and colour intensity being proportional to flow velocity, this is called Colour Doppler ultrasound. Spectral Doppler is a valuable tool to evaluate blood flow patterns quantitatively whereas Colour Doppler provides a more graphical depiction of vascular anatomy similar to an X-ray angiogram (Figure 1).

There is a third mode of presentation is a variation on the Colour Doppler technique called Power Doppler ultrasound, whereby the colour intensity does not reflect the mean flow velocity at each point in time but all flow velocities summated; this provides less indication about the character of the blood flow but is much more sensitive to small vessels and slow blood flow (Figure 2).


**Figure 2:** Power Doppler ultrasound is used to detect slow flow in small vessels; this image of the testis shows the presence of venous and arterial blood flow and therefore excludes the possibility of testicular torsion.




**Figure 3:** Normal triphasic pattern of peripheral arterial spectral Doppler: (a) corresponds to cardiac systole and (b) and (c) represent vascular recoil.



**Figure 4:** The pre-stenotic tardus parvus waveform results from dampening and delaying of flow caused by the resistance induced by an on-coming stenosis in this case in the hepatic artery.




(a) shows superior mesenteric artery stenosis




**Figure 5:** Flow acceleration occurs within a blood vessel at the site of stenosis with increased systolic and diastolic velocities:

(b) high grade superior mesenteric artery stenosis with flow velocities >260cm/s.



**Figure 6:** Post stenotic waveforms show increased systolic velocity but reduced or no diastolic velocity/flow; note that flow below the base line may not indicate reversed flow as it depends on the orientation of the probe angle to the direction of flow.



end diastolic flow velocities; when the arterioles are open end diastolic flow is high, while when they are closed it is low. Peak systolic flow is essentially constant in both states.

The factor most commonly used to assess arteriolar tone (and hence organ activity) is the Resistive Index, which is calculated as follows:

$$RI = (\text{peak systolic velocity} - \text{end diastolic velocity}) / \text{peak systolic velocity}$$

A high RI (>0.7) occurs when an organ is in “power save” mode, while a low RI (0.55-0.7) is seen when the organ is “on”. RIs outside of the range expected for the organ’s state usually indicate disease. In the hepatic artery, a high RI may indicate a post prandial state (ie patient did not fast) or diffuse distal microvascular disease as seen in chronic liver disease such as cirrhosis or chronic hepatitis. A low RI in the hepatic artery is more indicative of disease and may occur with proximal arterial stenosis or distal arteriovenous or arterioportal shunting as may occur in severe cirrhosis.

Other indices besides RI that are used to assess arteriolar tone are systolic-diastolic ratio (PSV/EDV) and pulsatility index ( $PI = (PSV - EDV) / V_{mean}$ ).

Central arteries have a monophasic pattern with only antegrade flow unlike peripheral arteries that show recoil. This is due to low arteriolar tone that ensures

### Arterial Spectral Waveforms

Spectral Doppler waveforms can be used to identify blood vessels (“vascular signatures”). Classical peripheral arterial flow has a triphasic pattern (Figure 3) with the first phase (largest wave) reflecting cardiac systole and the subsequent two phases correlating to healthy arterial wall recoil. Loss of the second and third phases resulting in a biphasic or monophasic pattern indicates loss of arterial wall elasticity that results from atherosclerosis.

Loss of wave amplitude and a delayed peak are the results of arterial stenosis located distally; this is referred to as a tardus parvus waveform (latin translation for late and low) (Figure 4). High wave amplitude means that there is acceleration of blood flow; this occurs at the site of vascular stenosis (Figure 5). This phenomenon can be reproduced by pinching the end of your garden hose; this will obtain a more powerful jet to transmit water to farther corners of your garden. Both peak systolic and end diastolic flow velocities are increased at the site of stenosis. Distal to an arterial stenosis, peak systolic flow velocity is still increased but to a lesser extent while end diastolic flow velocity is diminished or absent (Figure 6).


Distal to an arterial occlusion there is no flow; this is termed an aphasic

flow pattern (in distinction to mono-, di- and triphasic flow patterns described earlier.

In a normal healthy state, arterioles have the capacity to change their resistance to divert flow toward the organs that need it most. Organs that need to be “on” have their arterioles relaxed so that they are appropriately perfused. When an organ is on “power save” mode, its arterioles constrict and flow is diverted to other organs. The brain, the liver, the kidneys and the gonads need to be “on” at all times, so its arterioles are relaxed. During rest, skeletal muscles are on “power save” mode so their arterioles are constricted. The same can be said for intestinal vessels in a fasting state (which is the case during most abdominal ultrasound examinations). During active physical exercise, skeletal muscle arterioles relax to allow delivery of more oxygenated blood to those tissues. The same holds true for intestinal arterioles in the post-prandial state. Arteriolar tone can be analysed with spectral Doppler imaging by observing the relationship of peak systolic and

A persistently high hepatic arterial RI may be seen in patients of advanced age and diffuse peripheral microvascular (arteriolar) compression or disease


**Figure 7:** “Power-save” flow mode in the external artery




(a) with low diastolic flow velocity (\*) compared to “always on” flow mode in the internal carotid artery

(b) with high diastolic flow velocity (\*\*); consequently high RI in the ECA and low RI in the ICA.


**Figure 8:** Phasicity in peripheral venous blood flow is the result of changes in intrathoracic and intra-right atrial pressures.



**Figure 9:** Hepatic vein ASCD waveform.



**Figure 10:** Portal vein phasicity is low amplitude indirectly reflecting changes in central venous pressure.



16-40cm/s. Slow flow (<10cm/s) is seen in the portal vein in cases of cirrhosis with portal hypertension with no flow or flow reversal developing in severe cases. Note however that portal venous thrombosis (eg idiopathic or secondary to malignant disease) would also result in absent flow.

As is evident from the above data, Doppler ultrasound is a powerful tool that can help diagnose disease and also follow disease progression. Combining flow information with grey scale ultrasound findings can provide information that is unmatched by other imaging modalities. New flow encoding modalities and more sensitive and low interference technologies have made Doppler ultrasound an indispensable tool in the diagnostic imaging department. <sup>5</sup>

delivery of blood to central organs at all times. RI will however differ depending on organ requirements with the internal carotid and hepatic arteries having a lower RI (RI 0.55-0.7) than the external carotid and renal arteries (RI > 0.7) (Figure 7). RI values also change with organ state; the hepatic arterial RI in the fasting state is higher (>0.7) than that in the post prandial state (0.55-0.7).

A persistently high hepatic arterial RI may be seen in patients of advanced age and diffuse peripheral microvascular (arteriolar) compression or disease, as seen in chronic hepatocellular disease (including cirrhosis), hepatic venous congestion, cold ischemia (post-transplantation), and any stage of transplant rejection. Low hepatic arterial RI may occur with proximal arterial narrowing (transplant hepatic artery stenosis, atherosclerotic disease [celiac or hepatic], arcuate ligament syndrome) and distal (peripheral) vascular shunts (post-traumatic or iatrogenic arteriovenous fistulas, cirrhosis with portal hypertension and associated arteriovenous or arteriportal shunts, Osler-Weber-Rendu syndrome with arteriovenous fistulas).

### Venous Spectral Waveforms

Venous Spectral Waveforms normally show less pulsatility than arterial ones. Since the systemic venous system is returning blood to the heart, the flow direction is reversed compared to arterial flow. Also since blood is being transferred to the chest, changes in intrathoracic pressure will affect flow velocity with increased flow

during inspiration (due to negative intrathoracic pressure) (Figure 8).

When evaluating flow pattern in systemic veins that are closer to the heart (eg jugular and hepatic veins), changes in right atrial pressure also influence flow. Thus an ASCD waveform similar to that seen in the both jugular and hepatic veins (Figure 9 ASCD in hepatic veins) with an upward A wave corresponding to right atrial contraction, a downward S wave resulting from antro-ventricular septal traction caused by ventricular contraction that reduces intra-atrial pressure, a V wave that correlates with right atrial filling, and a D wave correlating with opening of the tricuspid valve. Thus an abnormal tricuspid valve (stenosis or regurgitation), right heart failure or a cardiac septal defect will result in abnormal hepatic venous flow waveforms.

The portal venous waveform differs from systemic ones in that changes in intrathoracic and right atrial pressures have less of an influence on flow due to the intervening liver. The normal portal venous waveform should gently undulate and always remain above the baseline. The peak portal velocity ( $V_1$ ) corresponds to systole, and the trough velocity ( $V_2$ ) corresponds to end diastole (Figure 10). Abnormally high pulsatility in the portal vein may be due to cardiac causes as is seen in tricuspid regurgitation and right-sided CHF or due to arteriovenous shunting (as seen in severe cirrhosis) or arteriovenous fistulas (as seen in hereditary hemorrhagic telangiectasia). Normal flow velocity in the portal vein is

