

# DETECTING LOW-LEVEL SEWAGE POLLUTION USING ROCKY SHORE COMMUNITIES AS BIO-INDICATORS

Adrian MALLJA & Patrick J. SCHEMBRI

Department of Biology, University of Malta, Msida, Malta.

While coastal pollution due to high inputs of organic matter is easy to detect and monitor, this is much more difficult in the case of sporadic low-level inputs. Moreover, routine water-quality surveys of large stretches of coastline are time-consuming and often prohibitively expensive. Such monitoring is therefore usually limited to sensitive areas. These restrictions make the results less useful for purposes of coastal pollution management. The indirect assessment of the degree of pollution is thus very appealing



Fig. 1. The Maltese Islands - the location of Xghajra and the 4 control sites relative to the sewage outfall.

(SATSMADHIS, 1985), more so when one can use inexpensive equipment and perennially present indicators. Rocky shore community structure has the potential of being a very suitable indicator of coastal low-level organic pollution: it represents the integrated response of the shore biota to environmental perturbations over time and such communities are readily accessible from the land. The present study evaluates the suitability of using rocky shore community structure as such an indicator in the Maltese Islands. The rocky shore communities at Xghajra, located 1.3 km south of Malta's main sewage outfall and downstream from it, and those at four control sites north of the outfall (Fig. 1), were sampled quantitatively by means of 0.5 m x 0.05 m contiguous quadrats along belt transects set perpendicular to the shoreline. Six transects were sampled at Xghajra and one each at the control sites. Faunal species were recorded as number of individuals per unit area and the algae as percentage cover. The data were subjected to a hierarchical cluster analysis using centroid linkage and the Bray-Curtis similarity coefficient for the quantitative data, and the Jaccard coefficient and centroid linkage for the presence/absence data (DIXON, 1988). This was done to correlate the groupings formed with environmental factors.

These statistical analyses gave similar results for all the transects, irrespective of the site. Quadrats from each transect were clustered into three distinct groups. The first group contained all the algae and most of the lower shore animals (including *Lepidochitona caerogata*, *Patella hispidipennis*, *Patella caerulea*, *Dendropoma petraeum*, etc.). This corresponds to the lower mesolittoral zone of PÉRES & PICARD (1964). The second group contained the barnacle *Chthamalus stellatus*, sometimes alone but more often together with one or more other species, such as *Littorina neritoides*, *Patella rustica*, *Monodonta turbinata*, coralline algae, cyanobacteria or terrestrial lichens. This corresponds to PÉRES & PICARD'S upper mesolittoral zone. The third and last group, corresponding to the supralittoral zone of PÉRES & PICARD, was composed of the upper shore quadrats with the gastropod *L. neritoides* either alone, as at Xghajra, or together with one or both of the barnacles *C. stellatus* and *C. depressus*. However, Xghajra differed from the control sites in having a higher species richness (Table 1), and a different suite of species (Fig. 2). In particular, Xghajra differed in having a near total absence of the *Cystoseira* cover found on other rocky shores in the Maltese Islands, with only a few stunted specimens of *C. stricta* and *C. compressa* recorded; the absence of species intolerant to pollution (e.g. *Padina pavonica*, *Aceretabularia arctifolium*); and the presence of a large number of pollution-tolerant species (e.g. *Pterocladia capillacea*, *Corallina elongata*, *Gelidium aciculate*, *Ulva rigida*, *Ectocarpus* spp. and *Cladophora* spp.).

Thus, while the general zonation patterns at Xghajra were similar to those of the four control sites, the shore community here exhibited some peculiarities when compared to the rest, especially in the type of species present and in their abundance. The dominant algae at Xghajra formed associations characteristic of environments having high organic loading in the water as shown in other parts of the Mediterranean and the Red Sea (CORMACI *et al.*, 1985; D'ANNA *et al.*, 1985; ISMAIL & AWAI, 1987; CORMACI & FURNARI, 1991). The presence at Xghajra of a large population of *Mylaster minimus*, a well known indicator of high nutrient levels (D'ANNA *et al.*, 1985), is indicative of high levels of nutrients in this locality. The chemical analyses carried out in this region confirm this (CHIRCOFF, 1992). The type of species, the species richness, their abundance, as well as their associations (especially those exhibited by the algae), at Xghajra, are unusual for Maltese rocky shores and to date have only been found in this area. These results suggest that rocky shore biotic assemblages may be useful indicators of low-level sewage pollution, at least under local conditions.

## REFERENCES:

- CHIRCOFF, P. (1992) An investigation on the sewage outfall at West Għajnsielem, Ġgantija II-Sea Dissolution Plan of Sewage, Univ. of Malta, 118pp.
- CORMACI, M. & FURNARI, G. (1991). Phytobenthic communities as monitor of the environmental conditions of the Rimosa coastline. *Ostrea* 17 suppl. 1, 117-130.
- CORMACI, M., FURNARI, G., GRACIOPPO, G., CULIOLINA, P. & MANNINO, A.M. (1985) Metodo sistematico per la valutazione degli aspetti inquinanti nella costa di Augusta (Siracusa) - Sicil. Arc. Geogr. Sci. Mer. 18(152): 829-850.
- D'ANNA, G., GRACIOPPO, G. & RIGGIO, S., 1985. Lineamenti biologici dei bacini di rifiuti di Balaterra (Sicilia occidentale). *Ostrea* 11: 389-399.
- DIXON, W. J., 1988 (ed.) *BMWP Statistical Software Manual*, Vol 1 & II. University of California Press, Berkeley.
- ISMAIL, N. S. & AWAI, J., 1987. Effects of sewage dumping on macrobenthic invertebrates in the Jordan Gulf of Aqaba, Red Sea. *Int. Revue ges. Hydrobiol.* 72 (2): 225-234.
- PÉRES, JM & PICARD, J., 1964. Nouveau manuel de botanique benthique de la mer Méditerranée. *Rev. Trav. Sta. mar. Endoume*, 18: 15-30.

Location	species richness	algae	animals
Xghajra	65	26	29
Harbour-Coghnaq	21	11	10
Gozo	27	16	11
Malta	33	22	11
Dotted-line-Xghajra	32	16	16

Table 1. Comparison of the species richness at Xghajra and the 4 control sites.

DIAGNOSA		CONTROL SITES (4 REPLICATES)
UPPER LITTORAL ZONE		
SPOROBLITTORAL ZONE		
	UPPER	
	LOWER	
	DIAGN	
	DIAGNO	
	DIAGNO	

Fig. 2. Comparison of zonation patterns at Xghajra and a generalized zonation pattern for the 4 control sites. \* denotes non-pollutous species or species commonly found in degraded or polluted situations.