

Trichopoda pennipes (Fabricius, 1781) - a new tachinid fly (Diptera) for Malta

Amante DARMANIN¹ & Pierfilippo CERRETTI²

Tachinidae is a diverse, cosmopolitan family of true flies currently with approximately 8,600 described species (O'HARA, 2013). Of these, about 900 have been found in Europe (PAPE *et al.*, 2015) and 61 species have been recorded for the Maltese archipelago (EBEJER, 2011). Tachinid larvae develop as endoparasitoids of other insects, although occasional records on other arthropods are also known (STIREMAN *et al.*, 2006; CERRETTI *et al.*, 2014). Four subfamilies are traditionally recognised, namely: Dexiinae, Exoristinae, Phasiinae and Tachininae. With the exception of a few species belonging to one tribe (Strongygastrini), the Phasiinae are composed of species which attack true bugs (Heteroptera). The North American native *Trichopoda pennipes* is a parasitoid of Pentatomidae (Stink bugs) and Coreidae (Leaf-footed bugs) such as the green stink bug (*Chinavia hilaris*), the squash bug (*Anasa tristis*), the cactus coreid (*Chelinidea vittiger*) and *Leptoglossus* spp. (ARNAUD, 1978). It is also known to parasitise the southern green stink bug (*Nezara viridula*) a species also known from the Maltese archipelago. The present note is to provide some information on *Trichopoda pennipes* which is here recorded for the first time from the Maltese Islands based on photographs taken by one of us (AD).

Trichopoda pennipes (Fabricius, 1781)

(Fig. 1)

Material examined: Malta, Sananard (limits of Zabbar), 29.viii.2016, 1 ♂, A. Darmanin.




Figure 1: *Trichopoda pennipes* – habitus photograph.

¹ 131, Lanata, Convent Street, Zabbar ZBR 1356, Malta. E-mail: amantedar@gmail.com

² Dipartimento di Biologia e Biotecnologie ‘Charles Darwin’, Sapienza Università di Roma. E-mail: pierfilippo.cerretti@uniroma1.it

Trichopoda pennipes was first recorded in Europe (Italy) by COLAZZA *et al.* (1996) based on a record from 1989. Later, based on a specimen collected in Tuscany and preserved at Museum La Specola in Florence, TSCHORSNIG *et al.* (2012) moved back the date of the arrival of *T. pennipes* to 1983. However, soon after its arrival, *T. pennipes* rapidly spread throughout Italy (Campania, Friuli-Venezia Giulia, Liguria, Lombardia, Lazio, Umbria, Veneto, Sardinia and Sicily) (COLAZZA & BIN, 1990; GIANGIULIANI *et al.*, 1994; SALERNO *et al.*, 2002; CERRETTI *et al.*, 2004; CERRETTI, 2010; CARGNUS *et al.*, 2011). In other parts of Europe, it has been recorded in Portugal, Spain, France, Holland and Slovenia (TSCHORSNIG *et al.*, 2012).

The reproductive strategy and the life cycle of this fly is well known, because of its effectiveness in natural pest control systems. The adult female lays one to several plano-convex, non-incubated eggs on a large nymph or adult bug. Eventually, however, only one larva survives per host. The larva bores its way into the bug and feeds on its body fluids for about two weeks. The larva eventually kills the bug and emerges from the posterior abdominal segments. The third instar of *T. pennipes* then drops to the ground and pupates in the soil. An adult fly emerges about two weeks later.

ACKNOWLEDGMENTS

We would like to thank Prof David Mifsud for all his help during the preparation of this work.

REFERENCES

- ARNAUD, P.H. JR (1978) A host-parasite catalogue of North American Tachinidae (Diptera). *United States Department of Agriculture. Miscellaneous Publication*, 1319: 1–860.
- CARGNUS, E., BUIAN, F.M. & ZANDIGIACOMO, P. (2011) Presenza di *Trichopoda pennipes* (Diptera, Tachinidae) nell'Italia nord-orientale. *Bollettino Soc. Naturalisti.*, 35: 123–130.
- CERRETTI, P. (2010). I tachinidi della fauna italiana (Diptera Tachinidae), con chiave interattiva dei generi ovest-paleartici. Volume I, 573 pp., Volume II, 339 pp. + CD-ROM. Cierre Edizioni, Verona.
- CERRETTI, P., O'HARA, J.E., WOOD, D.M., SHIMA, H., INCLAN, D.J. & STIREMAN J.O. (2014) Signal through the noise? Phylogeny of the Tachinidae (Diptera) as inferred from morphological evidence. *Systematic Entomology*, 39: 335–353.
- CERRETTI, P., WHITMORE, D., MASON, F. & VIGNA TAGLIANTI, A. (2004) Survey on the spatio-temporal distribution of tachinid flies – using Malaise traps (Diptera, Tachinidae) (pp. 229–256). In: CERRETTI, P., HARDERSEN, S., MASON, F., NARDI, G., TISATO, M. & ZAPPAROLI M. [eds.], *Invertebrati di una foresta della Pianura Padana, Bosco della Fontana*. Secondo contributo. *Conservazione Habitat Invertebrati*, 3. Cierre Grafica Editore, Verona: 303 pp.
- COLLAZZA, S. & BIN, F. (1990) I Pentatomidi ed i loro entomofagi associate alla soia in Italia centrale. *Informatore fitopatologico*, 40 (2): 38–42.
- COLLAZZA, S., GIANGIULIANI, G. & BIN, F. (1996) Fortuitous introduction and successful establishment of *Trichopoda pennipes* F.: adult parasitoid of *Nezara viridula* (L.). *Biological Control*, 6 (3): 409–411.
- EBEJER, M. (2011) The Tachinidae (Diptera) of the Maltese Islands. *Bulletin of the Entomological Society of Malta*, 4 : 59–72.
- GIANGIULIANI, G., COLLAZZA, S., LUCA, G. & FARINELLI, D. (1994) Efficacia del parassitoide *Trichopoda pennipes* F. nella localizzazione dell'ospite *Nezara viridula* (L.). *Atti XVII Congresso nazionale italiano di Entomologia*, Udine 13–18 Giugno 1994: 597–600.
- O'HARA, J.E. (2013) History of tachinid classification (Diptera, Tachinidae). *ZooKeys*, 316: 1–34. doi: 10.3897/zookeys.316.5132

- PAPE, T., BEUK, P., PONT, A.C., SHATALKIN, A.I., OZEROV, A.L., WOŹNICA, A.J., MERZ, B., BYSTROWSKI, C., RAPER, C., BERGSTRÖM, C., KEHLMAIER, C., CLEMENTS, D.K., GREATHEAD, D., KAMENEVA, E.P., NARTSHUK, E., PETERSEN, F.T., WEBER, G., BÄCHLI, G., GELLER-GRIMM, F., VAN DE WEYER, G., TSCHORSNIG, H-P., DE JONG, H., VAN ZUIJLEN, J-W., VAŇHARA, J., ROHÁČEK, J., ZIEGLER, J., MAJER, J., HŮRKA, K., HOLSTON, K., ROGNES, K., GREVE-JENSEN, L., MUNARI, L., DE MEYER, M., POLLET, M., SPEIGHT, M.C.D., EBEJER, M.J., MARTINEZ, M., CARLES-TOLRÁ, M., FÖLDVÁRI, M., CHVÁLA, M., BARTÁK, M., EVENHUIS, N.L., CHANDLER, P.J., CERRETTI, P., MEIER, R., ROZKOSNY, R., PRESCHER, S., GAIMARI, S.D., ZATWARNICKI, T., ZEEGERS, T., DIKOW, T., KORNEYEV, V.A., RICHTER, V.A., MICHELSEN, V., TANASIJTSHUK, V.N., MATHIS, W.N., HUBENOV, Z. & DE JONG, Y. (2015) Fauna Europaea: Diptera – Brachycera. *Biodiversity Data Journal*, 3: e4187. doi: 10.3897/BDJ.3.e4187
- SALERNO, G., COLAZZA, S. & BIN, F. (2002) *Nezara viridula* parasitism by the tachinid fly *Trichopoda pennipes* ten years after its accidental introduction into Italy from the New World. *BioControl*, 47 (6): 617–624.
- STIREMAN, J.O., O'HARA J.E. & WOOD D.M. (2006) Tachinidae: evolution, behavior, and ecology. *Annual Review of Entomology*, 51: 525–555.
- TSCHORSNIG H.P, CERRETTI P. & ZEEGERS T. (2012) Eight “alien” tachinids in Europe? *The Tachinid Times*, 25 (Feb. 2012).